



Contribution ID: 402

Type: Postar

## Investigating the effects of exotic $\alpha$ -clustered nuclear density profiles on medium anisotropy in p–O and p–C collisions at the LHC

One of the major motivations for the planned p–O and O–O collisions at the LHC is to explore the possibilities of small system collectivity. Such transverse collective expansion results in the appearance of long-range azimuthal correlation and is quantified via the coefficients,  $v_n$ , of Fourier expansion of the azimuthal momentum distribution of the final-state particles. These flow coefficients serve as the medium response to the initial spatial anisotropy and are sensitive to the density profile of the colliding nuclei. Light nuclei such as  $^{12}\text{C}$  and  $^{16}\text{O}$  are theorized to possess extra stability due to the presence of an  $\alpha$ -clustered arrangement of its nucleons. In this context, studies on ultra-relativistic p–A collisions involving  $^{12}\text{C}$  or  $^{16}\text{O}$  nuclei can serve a dual purpose: exploring small system collectivity along with investigating the effects of a clustered nuclear geometry on the medium anisotropy. With this motivation, for the first time, we study p–O and p–C collisions at  $\sqrt{s_{\text{NN}}} = 9.9\text{ TeV}$  through a multi phase transport model (AMPT) simulations. We attempt to explore how an initial  $\alpha$ -clustered nuclear structure of  $^{16}\text{O}$  and  $^{12}\text{C}$  influences the production yield, initial eccentricities and flow coefficients in the final state, in comparison to an unclustered density profile, Sum-Of-Gaussians (SOG). The flow coefficients are estimated via a two-particle Q-cumulant method.

The results show that  $\langle\epsilon_2\rangle$  varies with centrality in a unique manner for  $\alpha$ -clustered p–O and p–C collisions, similar to O–O collisions. However, the centrality dependence of  $\langle\epsilon_2\rangle$  and  $\langle\epsilon_3\rangle$  is not effectively carried forward to the final state  $v_2$  and  $v_3$ , owing to lesser participants. We also see that the dependence of  $v_2$  on centrality is much less in comparison to  $v_3$ , which is reflected in the  $v_3/v_2$ ,  $v_2/\langle\epsilon_2\rangle$  and  $v_3/\langle\epsilon_3\rangle$  ratios studied in our work. We notice that the  $\alpha$ -clustered case shows almost a flatter trend of  $v_3$  with centrality than the corresponding collisions with a SOG profile, possibly indicating the discretized internal structure of an  $\alpha$ -clustered nucleus. Thus, by probing the effects of the nuclear structure employing ultra-relativistic collisions, this work serves as a transport-model-based prediction for the upcoming p–O collisions in the LHC Run 3.

### Field of contribution

Phenomenology

**Author:** K R, Aswathy Menon (Indian Institute of Technology Indore (IN))

**Co-authors:** Mr PRASAD, Suraj (Indian Institute of Technology Indore (IN)); MALLICK, Neelkamal (University of Jyväskylä); SAHOO, Raghunath (Indian Institute of Technology Indore (IN))

**Presenter:** K R, Aswathy Menon (Indian Institute of Technology Indore (IN))

**Track Classification:** Heavy ion and QCD