


Dark Neutrino and Explanation of MiniBooNE Anomaly Under Siege

Based on arXiv:1812.08768



Carlos A. Argüelles



also starring




Matheus Hostert (Durham) and Yu-Dai Tsai (FNAL)

The back story: LSND


- LSND used an 800 MeV proton beam to produce pions which decay at rest.



- Search for antineutrino-electron appearance
- Observed 3.8 sigma excess of events



If interpreted as an appearance probability ...




$$P(\nu_{\mu} \rightarrow \nu_e) = \sin^2(2\theta) \sin^2\left(1.27 \frac{\Delta m^2 L}{E_{\nu}}\right)$$

- The preferred parameter space of the LSND anomaly was not compatible with other known mass differences.
- If this is due to a new neutrino mass state, then we should observe a similar signal at different E and L, but same L/E!

A. Aguilar-Arevalo et al. [LSND Collaboration]
Phys. Rev. D 64, 112007 (2001) [hep-ex/0104049].

MiniBooNE@FNAL: proposed to test the LSND anomaly


- Approximately same L/E, but ~ 15 x larger energy and baseline.
- Decay-in-flight pion source.
- Higher backgrounds than LSND, but more statistics!
- Neutrino and antineutrino mode available.



MiniBooNE experimental signatures

Three typical event signatures:



- Muon-neutrino CCQE produces sharp photon ring on PMTS,
- Electron-neutrino CCQE events produces fuzzy ring,
- Muon-neutrino NC can produce π^0 : two gammas \rightarrow two fuzzy rings.




Cannot distinguish between electrons and photons

Recent MiniBooNE excess of neutrino-electron-like events

- MB has reported an excess in neutrino and antineutrino channels.
- They claim that this excess is compatible with LSND. (Previous results show tension with LSND in neutrino mode)
- Excess has remained after doubling the data: not statistical in nature.
- It has a significance of 4.7 sigma.




Neutrino flux panorama



A. Diaz, CA, G.Collin, JM. Conrad, M. Shaevitz to appear
 very soon.

Sterile neutrino panorama



A. Diaz, CA, G.Collin, JM. Conrad, M. Shaevitz to appear
very soon.

So ... We have discover a new particle!?



If it's a “vanilla” eV-scale sterile neutrino

$$P_{\nu_e \rightarrow \nu_e} = 1 - 4(1 - |U_{e4}|^2)|U_{e4}|^2 \sin^2(1.27\Delta m_{41}^2 L/E)$$

$$P_{\nu_\mu \rightarrow \nu_e} = 4|U_{e4}|^2|U_{\mu4}|^2 \sin^2(1.27\Delta m_{41}^2 L/E)$$

$$P_{\nu_\mu \rightarrow \nu_\mu} = 1 - 4(1 - |U_{\mu4}|^2)|U_{\mu4}|^2 \sin^2(1.27\Delta m_{41}^2 L/E)$$

Oscillation probabilities among appearance and disappearance channels are related.



Need to look in other channels for further confirmation!

So ... We have discover a new particle!?



* R.I.P. grumpy cat.

App and Dis preference regions don't match!



From Collin et al 1602.00671, similar conclusions from other groups see Gariazzo et al. 1703.00860, and Dentler et al JHEP 1808 (2018)

Tension in the global data!!!

PG: parameter goodness-of-fit.
Larger is better. Small is bad.
Very small is very bad.

Analysis	$\chi^2_{\min, \text{global}}$	$\chi^2_{\min, \text{app}}$	$\Delta\chi^2_{\text{app}}$	$\chi^2_{\min, \text{disapp}}$	$\Delta\chi^2_{\text{disapp}}$	$\chi^2_{\text{PG}}/\text{dof}$	PG
Global	1120.9	79.1	11.9	1012.2	17.7	29.6/2	3.71×10^{-7}
Removing anomalous data sets							
w/o LSND	1099.2	86.8	12.8	1012.2	0.1	12.9/2	1.6×10^{-3}
w/o MiniBooNE	1012.2	40.7	8.3	947.2	16.1	24.4/2	5.2×10^{-6}
w/o reactors	925.1	79.1	12.2	833.8	8.1	20.3/2	3.8×10^{-5}
w/o gallium	1116.0	79.1	13.8	1003.1	20.1	33.9/2	4.4×10^{-8}
Removing constraints							
w/o IceCube	920.8	79.1	11.9	812.4	17.5	29.4/2	4.2×10^{-7}
w/o MINOS(+)	1052.1	79.1	15.6	948.6	8.94	24.5/2	4.7×10^{-6}
w/o MB disapp	1054.9	79.1	14.7	947.2	13.9	28.7/2	6.0×10^{-7}
w/o CDHS	1104.8	79.1	11.9	997.5	16.3	28.2/2	7.5×10^{-7}
Removing classes of data							
$\bar{\nu}_e$ dis vs app	628.6	79.1	0.8	542.9	5.8	6.6/2	3.6×10^{-2}
$\bar{\nu}_\mu$ dis vs app	564.7	79.1	12.0	468.9	4.7	16.7/2	2.3×10^{-4}
$\bar{\nu}_\mu$ dis + solar vs app	884.4	79.1	13.9	781.7	9.7	23.6/2	7.4×10^{-6}

Tension in the global data!!!

PG: parameter goodness-of-fit.
Larger is better. Small is bad.
Very small is very bad.

Analysis	$\chi^2_{\min, \text{global}}$	$\chi^2_{\min, \text{app}}$	$\Delta\chi^2_{\text{app}}$	$\chi^2_{\min, \text{disapp}}$	$\Delta\chi^2_{\text{disapp}}$	$\chi^2_{\text{PG}}/\text{dof}$	PG
Global	1120.9	79.1	11.9	1012.2	17.7	29.6/2	3.71×10^{-7}
Removing anomalous data sets							
w/o LSND	1099.2	86.8	12.8	1012.2	0.1	12.9/2	1.6×10^{-3}
w/o MiniBooNE	1012.2	40.7	8.3	947.2	16.1	24.4/2	5.2×10^{-6}
w/o reactors	925.1	79.1	12.2	833.8	8.1	20.3/2	3.8×10^{-5}
w/o CDHS	1104.8	79.1	11.9	997.5	16.3	28.2/2	7.5×10^{-7}
Removing classes of data							
$\bar{\nu}_e$ dis vs app	628.6	79.1	0.8	542.9	5.8	6.6/2	3.6×10^{-2}
$\bar{\nu}_\mu$ dis vs app	564.7	79.1	12.0	468.9	4.7	16.7/2	2.3×10^{-4}
$\bar{\nu}_\mu$ dis + solar vs app	884.4	79.1	13.9	781.7	9.7	23.6/2	7.4×10^{-6}

A vanilla 3+1 sterile neutrino fails to explain all the data!

This raises some more questions...

Do we understand all SM background/process well enough?

Are all the anomalies related? Or only some of them?

E.g., are LSND and MiniBooNE observing the same physics?

Since null results are not scrutinized as carefully as anomalous ones. Are all null results reliable?

Is there a significant signal of electron-neutrino disappearance (e.g. reactors)?

If the anomalies are confirmed as new physics, in what theories are they embedded?

Novel alternative explanations of the MiniBooNE anomaly

Assume that the SM gauge group is extended by a new U(1)', which mixes kinetically with the SM hypercharge

$$\mathcal{L}_{\text{kin}} \supset \frac{1}{4} \hat{Z}'_{\mu\nu} \hat{Z}'^{\mu\nu} + \frac{\sin \chi}{2} \hat{Z}'_{\mu\nu} \hat{B}^{\mu\nu} + \frac{m_{\hat{Z}'}^2}{2} \hat{Z}'^\mu \hat{Z}'_\mu$$



Also introduce a new SM-gauge singlet, charged under the new U(1)', which is allowed to mix with SM active neutrinos.

Bertuzzo et al. Phys. Rev. Lett. 121, 241801 (2018)



Ballett et al. arXiv:1808.02915

Interaction Lagrangian

$$\mathcal{L} \supset -eq_f c_W \chi \bar{f} \gamma^\mu f Z'_\mu$$




$$U_{\alpha 4}^* g' \bar{\nu}_\alpha \gamma^\mu P_L \nu_4 Z'_\mu + U_{\alpha 4}^* U_{\beta 4} g' \bar{\nu}_\alpha \gamma^\mu P_L \nu_\beta Z'_\mu$$



Producing the MiniBooNE signature: two implementations

$$\alpha_D \alpha_{\text{QED}} \varepsilon^2 |U_{\alpha 4}|^2$$



Small Mz' :


- Larger cross section
- Small Q^2 : more forward ν_4 , larger coherent to diffractive contributions

Large Mz' :

- Smaller cross section
- Larger Q^2 : more isotropic ν_4 production, more diffractive contribution

Producing the MiniBooNE signature


**Model from Bertuzzo et al.
Phys. Rev. Lett. 121, 241801 (2018)**



If $M_4 > M_{Z'}$, two body decay is the dominant decay channel.
 $M_4 \gtrsim 100$ MeV so the decay products are not so boosted in order to reproduce angular distribution.

Producing the MiniBooNE signature



**Model from Bertuzzo et al.
Phys. Rev. Lett. 121, 241801 (2018)**



$M_{Z'}$ should be light (< 60 MeV) so that the electron pair is collimated and can “fake” an electron-neutrino ring


Model by Bertuzzo et al. parameter space

Distributions at benchmark point




Producing the MiniBooNE signature

Model from Ballett et al.
arXiv:1808.02915; see also 1903.07589.







If $M_4 < M_{Z'}$, three body decay is the dominant decay channel.



$M_4 \sim > 300$ MeV: large too many high-energy events
 $M_4 \sim < 50$ MeV most events in lowest energy bin.

$M_{Z'} \sim < 1$ GeV: spectrum is too forward.

Model by Ballett et al. parameter space






Siege

noun

a military operation in which enemy forces surround a town or building, cutting off essential supplies, with the aim of compelling the surrender of those inside.



Statistical limitations

Unknown nuSM parameters

Backgrounds


Flux Uncertainties

Cross section uncertainties

Strategy

NOTE that I am NOT putting scales in this diagram!

Looking for novel neutrino interactions requires understanding of our SM neutrino interactions. This is tough!



Large cross=section


Small cross=section



Strategy

NOTE that I am NOT putting scales in this diagram!

Looking for novel neutrino interactions requires understanding of our SM neutrino interactions. This is tough!



Large cross-section

Small cross-section



Let's focus on the realization by Bertuzzo et al.

- Production cross section is dominated by coherent processes; i.e. little/no hadronic activity at vertex.
- Products angular distribution is broad at Booster beam energies, but less so at higher energy beams.
- Electron pair produced by Z' decay is very collimated.

How can we confirm or rule out a model like this?

*It will become apparent soon why I focus on this model realization model

Let's focus on the realization by Bertuzzo et al.


- Production cross section is dominated by coherent processes; i.e. little/no hadronic activity at vertex.
- Products angular distribution is broad at Booster energies, but less so at higher energy beams.
- Electron pair produced by Z' decay is very collimated.

How can we confirm or rule out a model like this?

We are going to use neutrino scattering data to look for evidence of this process

*It will become apparent soon why I focus on this model realization model

How big of a cross section are we talking about?



Here we use benchmark (BP) point parameters reported by Bertuzzo et al. 1807.09877.

Neutrino-electron scattering measurements


We have measured neutrino-electron scattering @:

- **LSND**
 - **TEXONO**
 - **Borexino**
 - **SuperK**
 - **MINERvA Low-Energy**
 - **CHARM-II**
- Too low energies for BSM case of interest**
- We will focus on these experiments.**

Will measure it very soon @ **MINERvA Medium-Energy, NOvA,**
and later @ **DUNE.**

Strategy

Electron-neutrino-like scattering search



*how much *leakage* onto the large angle depends on model parameter and neutrino energy.

Working around limited information!

By design at final cut level CHARM-II and Minerva measurements have small backgrounds: also means small amount of BSM-signal leaking in. We cannot use the final event samples to constrain the new models :(

Would be great if we had access to the reconstructed electron energy and angular distributions at different cut levels.




Background,
e.g. NC, π^0



Neutrino-
electron
scat.



Recent BSM-MB
explanations





MINERvA analysis strategy

For MINERvA we are going to use the dE/dX distribution of candidate electron-neutrino scattering events.

All MINERvA cuts applied, except for the final dE/dX cut!

Note that backgrounds have been tuned here!





MINERvA analysis strategy

Parameter	Tuned value
ν_e	0.76 ± 0.03
ν_μ NC	0.64 ± 0.03
ν_μ CC	1.00 ± 0.02

For MINERvA we are going to use the dE/dX distribution of candidate electron-neutrino scattering events.

All MINERvA cuts applied, except for the final dE/dX cut!



Note that backgrounds have been tuned here!



Sidebands used for tuning background on MINERvA

Tuning parameters diagram from J. Park thesis:

<http://lss.fnal.gov/archive/thesis/2000/fermilab-thesis-2013-36.pdf>



For large heavy neutrino masses the BSM contribution leaks the sideband used to constrain the background on the neutrino electron scattering region.

MINERvA: Our Analysis setup

We use the following χ^2 definition:

$$\chi_{\alpha\beta}^2 = \frac{(N_{\text{data}} - (1 + \alpha + \beta)\mu_{\text{MC}}^{\text{BKG}} - (1 + \alpha)\mu_{\text{MC}}^{nu-e} - (1 + \alpha)\mu_{\text{BSM}})^2}{N_{\text{data}}} + \left(\frac{\alpha}{\sigma_{\alpha}}\right)^2 + \left(\frac{\beta}{\sigma_{\beta}}\right)^2$$


- We set $\sigma_{\alpha} = 10\%$ account for beam uncertainties.
- We set $\sigma_{\beta} = 30\%$ motivated by the amount of tuning; conservative with respect to tune normalization uncertainty.
- We include only coherent contribution to the BSM signal to avoid hadronic activity cuts.

MINERvA: Our Analysis setup


- We do a rate-only analysis on the single bin with

$dE/dx > 4.5 \text{ MeV}/(1.7 \text{ cm})$

- We use 3.43×10^{20} POTs, Assume fiducial mass of 6.10 tons.




MINERvA result




MINERvA result

We checked that changing the background uncertainty from 30% to 100% changes the result by no more than a factor of two. The constraint power is coming from the BSM signal overshooting the data.




CHARM-II: complementary measurement




For CHARM-II we are going to use the $E\theta^2$ distribution before the final dE/dX cut is applied.


Finding “BSM-safe” sideband to measure background




Angular distribution of BSM-signal



CHARM-II: complementary measurement



For CHARM-II we use the distribution before the angular cut and dE/dX were applied



Use the region with $E\theta^2 > 0.03$ to obtain the background uncertainty.


Allow for rate/slope to change; with this we estimate its rate to be constrained to be $\sim 3\%$.

Our CHARM-II analysis setup details

- Rate-only analysis on a single bin with $E \cos^2 \theta < 0.03$ GeV.
- Same χ^2 definition as in MINERvA, but updated uncertainties.
- Background norm. from sideband $\sim 3\%$; flux uncertainty $\sim 4\%$.
- We assume a fiducial mass of 547tons, $\langle A \rangle \sim 20.7$, and $2.5e19$ POT.

Putting it all together: the money plot

Here for CHARM-II we also consider 3 times larger background uncertainty (dashed)




Take home message: lessons learned

- We are excited to see upcoming neutrino-electron scattering analyses by Minerva-ME and NOvA!
- We have used two different experiments to constrain recent MiniBooNE explanations. Tensions are large with the realization given by Bertuzzo et al.
- These constraints are not effective for Ballett et al. due to the fact that coherent contributions are much smaller; thus the signal is not present due to hadronic activity cuts.
- Neutrino electron scattering is a powerful tool to constraint new physics interactions. Our constraints can also be adapted to other MB explanations, such as the dipole portal.


Bonus
slides!

MiniBooNE previous results

- Neutrino mode excess above the null primarily under 475 MeV. This is not expected given the LSND signal.
- Antineutrino mode saw excess above and below 475 MeV.
- Antineutrino mode was consistent with LSND, neutrino mode had tension.




Booster beam




- Maxima < 1 GeV neutrino energy
- Production of heavy states via neutrino interaction: hard.
- Heavier BSM physics look like “effective”-interactions; then angular distribution of excess wont work.

Recent proposals:
Light new physics $\sim < \text{GeV}$.


Dark Photon Searches





Heavy Neutrino Limits



See arXiv:1511.00683





$$|0\rangle = |\mathbf{v}_1\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad \text{and} \quad |1\rangle = |\mathbf{v}_2\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$