


# Quarkophobic $W'$ for LHC searches

Alfredo Gurrola - José Ruiz



# An initial motivation coming from $R(D^*)$


Low  
energy




High  
energy

A. Greljo, J. Camalich, J. Ruiz-Álvarez:  
Phys.Rev.Lett. 122 (2019) 13, 131803

# Competitive and complimentary





EFT




$W'$

# Quarkophobic $W'$



# Vector Boson Fusion $W'$ production at the LHC



# A simplified model implementation

1. Simplified: No complete model, only inclusion of vertices of interest.
2. Mimicking of SM TGC.
3. Minimal couplings: Only including what necessary to get processes.

$$\mathcal{L}_{VWW'}^1 = g_1^V V^\mu (W_{\mu\nu}^- W'^{+\nu} - W_{\mu\nu}^+ W'^{-\nu} + W_{\mu\nu}' W^{+\nu} - W_{\mu\nu}' W^{-\nu})$$


$$\mathcal{L}_{VWW'}^2 = g_2^V (W_\mu^+ W_\nu'^- V^{\mu\nu} + W_\mu'^+ W_\nu^- V^{\mu\nu})$$

$$V_{\mu\nu} = \partial_\mu V_\nu - \partial_\nu V_\mu \text{ and } V = Z \text{ or } \gamma$$


$$\mathcal{L}_l = \sum_l \bar{\nu}_l \gamma_\mu (g_l^R (1 + \gamma^5) + g_l^L (1 - \gamma^5)) W'^{\mu l}$$

# With the model implemented

1. Production of MC events:
  - a. MadGraph
  - b. Pythia 8
  - c. Delphes
2. Design a selection of events
3. Find exclusion limits
4. Possible caveats



# Differences among couplings







# Vector Boson Fusion $W'$ production at the LHC

Characteristics of the final state:

1. Two opposite hemispheres jets.
2. Large  $\eta$  separation among jets.
3. Large dijet invariant mass.
4. High  $p_T$  lepton.
5. High MET.



# Search designed for 1 TeV $W'$ (Z type 1 coupling)




$e$


$\mu$

$\tau$


# VBF variables



e




$\mu$




$\tau$


# VBF variables



e



$\mu$



$\tau$

# Selections

At least two jets with  $p_T > 60$  GeV,  $N(b)=0$ ,  $N(l)=1$ , other leptons veto  $p_T > 30$  GeV

Cut 1:  $\eta(j_1) \times \eta(j_2) < 0$

Cut 2:  $m_{jj} > 1000$  GeV

Cut 3:  $|\Delta(\eta(j_1), \eta(j_2))| > 4.0$

Cut 4:  $p_T(l) > 200$  GeV

Cut 5:  $p_T^{miss} > 200$  GeV

Cut 6:  $|\Delta(\phi(l), p_T^{miss})| > 1.0$

Cut 1:  $\eta(j_1) \times \eta(j_2) < 0$

Cut 2:  $m_{jj} > 1000$  GeV

Cut 3:  $|\Delta(\eta(j_1), \eta(j_2))| > 4.0$

Cut 4:  $p_T(l) > 150$  GeV

Cut 5:  $p_T^{miss} > 50$  GeV


Cut 6:  $|\Delta(\phi(l), p_T^{miss})| > 1.5$

e


$\mu$

$\tau$


# W' variables



e




$\mu$




$\tau$


# W' variables



e




$\mu$




$\tau$


# W' variables



e



μ



τ



# Cut flow: Electron and Muon

	Signal	W+jets	$t\bar{t}$	VV	$S/\sqrt{S+B}$		Signal	W+jets	$t\bar{t}$	VV	$S/\sqrt{S+B}$
Initial	1026.0	119107.0	36528.8	1453.0	2.6	Initial	1259.4	152156.8	48246.6	1870.3	2.8
Cut 1	751.8	53656.2	10932.3	609.1	2.9	Cut 1	930.3	68480.3	14364.6	788.2	3.2
Cut 2	487.5	10633.7	1353.6	112.2	4.3	Cut 2	593.7	13461.8	1796.9	147.8	4.7
Cut 3	427.5	2152.9	283.0	22.4	8.0	Cut 3	527.1	2771.8	365.0	27.9	8.7
Cut 4	380.7	93.9	11.2	1.8	17.2	Cut 4	474.0	118.4	17.6	2.1	19.2
Cut 5	371.1	75.1	9.4	1.4	17.4	Cut 5	459.9	91.7	11.6	1.9	19.3
Cut 6	370.2	0.0	0.0	0.7	19.2	Cut 6	459.6	0.0	0.0	0.9	21.4

e


$\mu$

## Cut flow: Tau


	Signal	W+jets	$t\bar{t}$	VV	$S/\sqrt{S+B}$
Initial	468.6	46671.3	17559.7	451.1	1.8
Cut 1	334.2	20778.8	5339.8	163.8	2.0
Cut 2	207.3	4085.0	703.3	26.5	2.9
Cut 3	178.5	860.5	137.0	5.0	5.2
Cut 4	145.5	129.5	36.6	0.8	8.2
Cut 5	145.5	129.5	36.6	0.8	8.2
Cut 6	136.8	53.6	15.7	0.3	9.5

$\tau$


# W' variables



e



$\mu$



$\tau$

# Some further considerations

$g_{VWW}$

$R(D^*)$

LHC

?

LHC

$M(W')$  [GeV]

# Conclusions

1. TGC with a  $W'$  give a new unexplored signature at the LHC.
2. TGC with a  $W'$  might be interesting for  $R(D^*)$  anomalies.
3. Implemented a simplified  $W'$  model with TGC for LHC searches.
4. Designed a search at the LHC for VBF produced high mass  $W'$ .
5. Proven sensitivity in this search.
6. Tau channel certainly more challenging than other leptons.
7. Low mass  $W'$  will be, at least, very challenging at the LHC.

**Thanks**

# Efficiencies: Electron and Muon

	Signal [%]	W-jets [%]	$t\bar{t}$ [%]	VV [%]		Signal [%]	W-jets [%]	$t\bar{t}$ [%]	VV [%]
Cut 1	73.27	45.05	29.93	41.92	Cut 1	73.87	45.01	29.77	42.14
Cut 2	47.51	8.93	3.71	7.72	Cut 2	47.14	8.85	3.72	7.9
Cut 3	41.67	1.81	0.77	1.54	Cut 3	41.85	1.82	0.76	1.49
Cut 4	37.11	0.08	0.03	0.12	Cut 4	37.64	0.08	0.04	0.11
Cut 5	36.17	0.06	0.03	0.10	Cut 5	36.52	0.06	0.02	0.1
Cut 6	36.08	<0.06	<0.03	0.05	Cut 6	36.49	<0.06	<0.02	0.05

e

$\mu$

## Efficiencies: Tau

	Signal [%]	W+jets [%]	$t\bar{t}$ [%]	VV [%]
Cut 1	71.32	44.52	30.41	36.3
Cut 2	44.24	8.75	4.0	5.87
Cut 3	38.09	1.84	0.78	1.12
Cut 4	31.05	0.28	0.21	0.17
Cut 5	31.05	0.28	0.21	0.17
Cut 6	29.19	0.11	0.09	0.07

$\tau$