



Contribution ID: 32

Type: **Parallel talk**

Looking for lepton-number-violating processes in $|\Delta L| = 2$ decays of B_s meson and Λ_b baryon

Friday 7 December 2018 11:30 (20 minutes)

Lepton-number violation can be induced by the exchange of an on-shell Majorana neutrino N in rare semileptonic $|\Delta L| = 2$ decays of the B_s meson and Λ_b baryon. We investigate the production of such a heavy sterile neutrino through these four-body $\mu^+\mu^+$ channels and explore the sensitivity that can be reached at the LHCb and CMS experiments. For heavy neutrino lifetimes of $\tau_N = [1, 100, 1000]$ ps and integrated luminosities collected of 10 and 50 fb^{-1} at the LHCb and 30, 300, and 3000 fb^{-1} at the CMS, we find a significant sensitivity on branching fractions of the orders $\sim 10^{-9} - 10^{-8}$. In the kinematically allowed mass ranges of m_N , we exclude regions on the parameter space $(m_N, |V_{\mu N}|^2)$ associated with the heavy neutrino.

Author: QUINTERO, Nestor (Universidad Santiago de Cali)

Co-authors: MILANES, Diego; MEJIA-GUISAO, Jhovanny; RUIZ-ALVAREZ, Jose

Presenter: QUINTERO, Nestor (Universidad Santiago de Cali)