



Contribution ID: 20

Type: Poster

## Low Energy Theorems and the Higgs boson.

Wednesday 4 December 2019 16:25 (5 minutes)

Decayings that have a lepton flavor violation (LFV) are affected by the standard model (SM). However, from recent observations at the LHC, some limits have been found for these types of processes that show a change in flavor, up to a confidence level of  $3\sigma$ . In particular, the process  $h \rightarrow \tau\tau$  seems to maintain a higher decay ratio (branching ratio)  $\mathcal{B}(h \rightarrow \tau\tau) \approx 0.25$ , compared to other channels. In this work we study the width of decay  $\Gamma(h \rightarrow \tau\tau)$  and  $\mathcal{B}(h \rightarrow \tau\tau)$  for this decay. The specification of both parameters is made using the following premises: The theoretical framework is that of the SM, but considers the neutrinos with mass, Both, 't Hooft - Feynman, and Unitary gauges, are used in the calculation. The Low Energy Theorems (LETs) technique is used to make the calculus to a one loop approach. In the end, the result obtained is contrasted with other results reported in the literature that have been performed using some other calculation techniques.

**Author:** GARCÍA, Roberto (BUAP)

**Presenter:** GARCÍA, Roberto (BUAP)

**Session Classification:** Poster session