

# Baryogenesis in the Standard Model EFT with dim 6 terms

Marta Losada  
New York University Abu Dhabi

Work done in collaboration with E. Fuchs, Y. Nir and Y. Viernik

Cosmology in Colombia  
Sept 25 2020

# Motivation I

- Observation of asymmetry of matter-antimatter in the Universe.

$$Y_B^{\text{obs}} = (8.59 \pm 0.08) \times 10^{-11}$$

Planck

- *How to make a matter filled Universe?*

- *Baryon number violation*
- *Departure from thermal equilibrium*
- *C and CP violation*

Sakharov '67

# Motivation II

- High precision, low energy experiments that have strong bounds.
- New CP violating terms have implications in other types of observables such as electric dipole moments.
- ACME bound

$$|d_e^{\max}| = 1.1 \times 10^{-29} \text{ e cm at 90% C. L.}$$

# Motivation III

- LHC results of Higgs discovery and measurement of physical properties.
  - Scalar particle with SM Higgs boson properties
- No new elementary particles discovered with  $m \sim 1 \text{ TeV}$ 
  - New physics scale is high enough to be parametrized via higher dimensional operators.
- New physics via higher dimension operators
  - Use Higgs physics results from LHC to constrain these higher dim terms
  - Consider dimension six terms of Higgs- fermion fields with complex couplings

# SM EFT Framework

- Dim-6 term with real and imaginary Yukawa in Lagrangian:

$$\mathcal{L}_{\text{Yuk}} = y_f \overline{F_L} F_R H + \frac{1}{\Lambda^2} (X_R^f + i X_I^f) |H|^2 \overline{F_L} F_R H + \text{h.c.}$$

- Allows for new CPV interactions
- Changes the fermion mass and the corresponding Yukawa coupling relation.

Parametrize by ratio of dim 6 to dim 4 contribution to the fermion mass:

$$T_R^f \equiv \frac{v^2}{2\Lambda^2} \frac{X_R^f}{y_f}, \quad T_I^f \equiv \frac{v^2}{2\Lambda^2} \frac{X_I^f}{y_f}$$

$$H = \frac{1}{\sqrt{2}}(v + h)$$

# Basics of EWBG

Kuzmin, Rubakov, Shaposhnikov '85

- Initial hot plasma with zero net baryon number with EW symmetry.
- As Universe expands and cools until EWPT around  $T \sim 100$  GeV
- Bubbles of the broken phase nucleate and expand to fill the Universe.
- Necessary to have new physics
  - CP violation sources: plasma particles CPV interactions with the bubble wall
  - Strong first order phase transition: suppress sphaleron transitions in the broken phase

# Electroweak Phase Transition

Assume:

- New degrees of freedom that produce a strong first order EWPT
- These do not affect the CPV interactions with bubble wall we are going to consider.
- No new sources of CPV from these new degrees of freedom.

There are important parameters such as the wall velocity and wall width that need to be obtained in a specific model. We will simply take on some benchmark values for them in this analysis.

# SMEFT implications for EW Baryogenesis

Full dynamics given by set of coupled equations:

$$\partial_\mu f^\mu = -\Gamma_M^f \mu_M^f - \Gamma_Y^f \mu_Y^f + \Gamma_{ss}^f \mu_{ss}^f - \Gamma_{ws}^f \mu_{ws}^f + S_f$$

$$\partial_\mu f^\mu \approx v_w f' - D_f f''$$


↗  
wall velocity

↗  
Diffusion constant

↑  
CPV Source term

# Source, relaxation and Yukawa terms

Use vev-insertion approx.



Lee et al

$$S_f \propto \text{Im}(m_f^* m'_f) \propto y_f^2 T_I^f$$

# Changes in CP-even rates

$$\Gamma_M \rightarrow \left[ \frac{(1 + r_{N0}^2 T_R^f)^2 + (r_{N0}^2 T_I^f)^2}{(1 + T_R^f)^2 + T_I^{f2}} \right] \Gamma_M$$

$$\Gamma_Y \rightarrow \left[ \frac{(1 + 3r_{N0}^2 T_R^f)^2 + (3r_{N0}^2 T_I^f)^2}{(1 + T_R^f)^2 + T_I^{f2}} \right] \Gamma_Y$$

# Baryon Asymmetry of the Universe

Using benchmark parameters


$$T_R^f = 0$$

$$Y_B = 8.6 \times 10^{-11} \times (51 T_I^t - 23 T_I^\tau - 0.44 T_I^b)$$

BAU for values:

$$|T_I^t| = \mathcal{O}(0.02), \quad |T_I^\tau| = \mathcal{O}(0.04), \quad |T_I^b| > 1$$

# SMEFT implications for (e)-EDMs



ACME bound

$$|d_e^{\max}| = 1.1 \times 10^{-29} \text{ e cm at 90% C. L.}$$

$$\frac{d_e^{(\textcolor{blue}{t})}}{e} \simeq -\frac{32\sqrt{2}}{3} \frac{e^2}{(16\pi^2)^2} \frac{m_e}{v^2} \left[ \left( 2 + \ln \frac{m_t^2}{m_h^2} \right) \left( \frac{y_t}{y_t^{\text{SM}}} \right)^2 T_I^t \right]$$

Panico et al '19

$$\frac{d_e^{(\textcolor{green}{b})}}{e} \simeq -\frac{32\sqrt{2}}{3} \frac{e^2}{(16\pi^2)^2} \frac{m_e}{v^2} \left[ \frac{1}{4} \left( \frac{\pi^2}{3} + \ln^2 \frac{m_b^2}{m_h^2} \right) \frac{m_b^2}{m_h^2} \left( \frac{y_b}{y_b^{\text{SM}}} \right)^2 T_I^b \right]$$

$$\frac{d_e^{(\tau,\mu)}}{e} \simeq -\frac{32\sqrt{2}}{3} \frac{e^2}{(16\pi^2)^2} \frac{m_e}{v^2} \left[ \frac{3}{4} \left( \frac{\pi^2}{3} + \ln^2 \frac{m_{\tau,\mu}^2}{m_h^2} \right) \frac{m_{\tau,\mu}^2}{m_h^2} \left( \frac{y_{\tau,\mu}}{y_{\tau,\mu}^{\text{SM}}} \right)^2 T_I^{\tau,\mu} \right]$$

# e-EDMs of only third generation fermions


$$d_e \approx |d_e^{\max}| \left[ 2223 \left( \frac{y_t}{y_t^{\text{SM}}} \right)^2 T_I^t + 9.6 \left( \frac{y_\tau}{y_\tau^{\text{SM}}} \right)^2 T_I^\tau + 11.6 \left( \frac{y_b}{y_b^{\text{SM}}} \right)^2 T_I^b \right]$$

So for,  $y_f = \mathcal{O}(y_f^{\text{SM}})$

$$T_I^t = \mathcal{O}(0.0004), \quad T_I^\tau = \mathcal{O}(0.1), \quad T_I^b = \mathcal{O}(0.09)$$

# SMEFT implications for Colliders

- Modification of Higgs production and decay modes.



$$\mu_I^F \equiv \frac{\sigma_I(pp \rightarrow h) \cdot \Gamma(h \rightarrow F)/\Gamma_h}{[\sigma_I(pp \rightarrow h) \cdot \Gamma(h \rightarrow F)/\Gamma_h]_{\text{SM}}}$$

$$r_f \equiv \frac{|\lambda_f|^2 / |\lambda_f^{\text{SM}}|^2}{|m_f|^2 / |m_f^{\text{SM}}|^2} = \frac{(1 + 3T_R^f)^2 + 9T_I^{f2}}{(1 + T_R^f)^2 + T_I^{f2}}$$

# Modified Production Rates, Decays and Total Width

Production Rates

$$\sigma_{\text{ggF}}/\sigma_{\text{ggF}}^{\text{SM}} = \sigma_{t\bar{t}h}/\sigma_{t\bar{t}h}^{\text{SM}} = r_t$$

$$\sigma_{Vh}/\sigma_{Vh}^{\text{SM}} = \sigma_{\text{VBF}}/\sigma_{\text{VBF}}^{\text{SM}} = 1$$

Decay Rates

$$\Gamma(h \rightarrow f\bar{f})/[\Gamma(h \rightarrow f\bar{f})]^{\text{SM}} = r_f \quad (f = b, \tau, \mu)$$

Total Width

$$\Gamma_h/\Gamma_h^{\text{SM}} = 1 + \text{BR}_b^{\text{SM}}(r_b - 1) + \text{BR}_\tau^{\text{SM}}(r_\tau - 1) + \text{BR}_g^{\text{SM}}(r_t - 1)$$

# LHC Measurements

channel	experiment	$\sqrt{s}$ / TeV	$\mathcal{L}/\text{fb}^{-1}$	comment	$\mu$
$h \rightarrow \tau^+ \tau^-$	ATLAS+CMS	7+8	5 + 20		$1.11^{+0.24}_{-0.22}$
	ATLAS	13	36.1	ggF, VBF	$1.09^{+0.35}_{-0.30}$
	CMS	13	77	ggF, $\bar{b}b$ , VBF, $Vh$	$0.75 \pm 0.17$
	ATLAS+CMS	7+8+13		all prod., priv. comb.	$0.91 \pm 0.13$
$h \rightarrow \mu^+ \mu^-$	ATLAS	13	139	upper bound at 95% C.L.	$< 1.7$
	CMS		35.9		$< 2.9$
$h \rightarrow \bar{b}b$	ATLAS	13	79.8	VBF+ $VH$ $t\bar{t}h + th$	$1.23 \pm 0.26$ $0.79^{+0.60}_{-0.59}$
	CMS	7+8+13	41.3	VH (0-2 $\ell$ , 2 b-tags+jets) all prod.	$1.01 \pm 0.22$ $1.04 \pm 0.2$
	ATLAS+CMS	7+8+13		VH, priv. comb. all prod., priv. comb.	$0.98 \pm 0.15$ $1.02 \pm 0.14$

+ all processes with t


# Combined flavors

$\sigma_I$	$\Gamma(h \rightarrow F)$	$\Gamma_h$	$f_1, f_2$	process	dependence
SM	$f_1$	$f_1, f_2$	$\tau, b$ $t, \tau$ $t, b$	any production, $h \rightarrow \tau\tau, b\bar{b}$ $Vh + \text{VBF}, h \rightarrow \tau\tau$ $Vh + \text{VBF}, h \rightarrow b\bar{b}$	A
$f_1$	SM	$f_1, f_2$	$t, b/\tau$	$ggF + tth, h \rightarrow VV$	
$f_1$	$f_2$	$f_1, f_2$	$t, \tau$ $t, b$	$ggF + tth, h \rightarrow \tau\tau$ $ggF + tth, h \rightarrow b\bar{b}$	

A: 
$$\mu_{\text{SM}}^{f_1} = \mu_{f_1}^{\text{SM}} = \frac{r_{f_1}}{\Gamma_h / \Gamma_h^{\text{SM}}} = \frac{r_{f_1}}{1 + \text{BR}_{f_1}^{\text{SM}}(r_{f_1} - 1) + \text{BR}_{f_2}^{\text{SM}}(r_{f_2} - 1)}$$

B: 
$$\mu_{f_1}^{f_2} = \frac{r_{f_1} r_{f_2}}{\Gamma_h / \Gamma_h^{\text{SM}}} = \frac{r_{f_1} r_{f_2}}{1 + \text{BR}_{f_1}^{\text{SM}}(r_{f_1} - 1) + \text{BR}_{f_2}^{\text{SM}}(r_{f_2} - 1)}$$

# Third generation quarks --bottom




All production modes combined

$$\mu_{b\bar{b}} = 1.02 \pm 0.14$$

dominated by

$$\mu_{Vh}^{bb}$$

# Third generation quarks –top (zoomed)



Constrained by  $\mu_{\text{ggF}}$ ,  $\mu_{t\bar{t}h}$  and  $\mu_{\gamma\gamma}$

$$\mu_{\text{ggF}+t\bar{t}h} = 1.09 \pm 0.08$$

# Leptons --muon

$$\frac{Y_B^{(\mu)}}{8.6 \times 10^{-11}} = \frac{d_e^{(\mu)}}{4.1 \times 10^{-30} e \text{ cm}}$$

$$\mu_{\mu^+\mu^-} = \frac{\Gamma(h \rightarrow \mu^+\mu^-)}{[\Gamma(h \rightarrow \mu^+\mu^-)]_{\text{SM}}}$$

$$\mu_{\mu^+\mu^-} = \frac{(1 + 3T_R^\mu)^2 + 9T_I^{\mu 2}}{(1 + T_R^\mu)^2 + T_I^{\mu 2}}$$

$\mu_{\mu^+\mu^-}^{\text{CMS}} < 2.9$  at 95% C.L.


$\mu_{\mu^+\mu^-}^{\text{ATLAS}} < 1.7$  at 95% C.L.

# Leptons --muon

$$\mu_{\mu^+\mu^-} = \frac{\Gamma(h \rightarrow \mu^+\mu^-)}{[\Gamma(h \rightarrow \mu^+\mu^-)]_{\text{SM}}}$$

$\mu_{\mu^+\mu^-}^{\text{CMS}} < 2.9$  at 95% C.L.

$\mu_{\mu^+\mu^-}^{\text{ATLAS}} < 1.7$  at 95% C.L.




$$|Y_B^{(\mu)}|_{\max} = 1.4 \times 10^{-11}$$
$$|d_e^{(\mu)}|_{\max} = 6.5 \times 10^{-31} \text{ e cm}$$

# Leptons- tau

$\Gamma(h \rightarrow \tau^+ \tau^-)$  and  $\Gamma_h$


All production modes

$$\mu_{\tau^+ \tau^-} = 0.91 \pm 0.13$$




# Combined Sources t-b

$$Y_B^{(t+b)} \lesssim 0.12 Y_B^{\text{obs}}$$



$$\begin{aligned} \mu_{t\bar{t}h+ggF}^{b\bar{b}} &= 0.88 \pm 0.43 \\ \mu_{VH}^{bb} &= 0.98 \pm 0.15 \\ \mu_{\text{ggF}+tth}^{VV} &= 1.08 \pm 0.08 \end{aligned}$$



# Combined Sources tau- b


No modification to production rates

$$\begin{aligned} \mu_{\tau^+\tau^-} \\ \mu_{b\bar{b}} \end{aligned}$$



Only Constraints

$$Y_B^{b+\tau,\max}(T_I^\tau = -0.4, T_I^b = +0.4) \simeq 7.8 Y_B^{\text{obs}}$$

$$T_R^f = 0$$



# Combined sources tau-t



$T_R^f = 0$

$$Y_B^{t+\tau, \text{max}} = Y_B^{t+\tau}(T_l^\tau = -0.3, T_l^t = +0.0016) \simeq 6.4 Y_B^{\text{obs}}$$

$$\mu_{\text{ggF}}^{\tau\tau} = 0.99 \pm 0.44$$

$$\mu_{\text{VBF+Vh}}^{\tau\tau} = 1.09 \pm 0.26$$

$$\mu_{\text{ggF+}t\bar{t}\text{h}}^{VV} = 1.08 \pm 0.08$$

SM-like solutions

$$Y_B = Y_B^{\text{obs}} \text{ with } d_e \simeq 0 \text{ and } \mu_f^F \simeq 1$$

Choose

$T_I^\tau$  and  $T_I^b$  such that  $d_e = 0$ ,

$T_R^\tau$  and  $T_R^b$  such that  $\mu_b = \mu_\tau = 1$

$$Y_B^{b+\tau, \text{max}}(d_e = 0, \mu_b = \mu_\tau = 1) = 10.25 Y_B^{\text{obs}}$$

# Implication for EFT Scales

Upper bounds on  $T_{I,R}$  from collider and EDMs.

$$\Lambda/\sqrt{X_{R,I}^f} \gtrsim \frac{\nu}{\sqrt{2}} \frac{1}{(y_f T_{R,I})^{1/2}} \sim \text{few} - \mathcal{O}(10) \text{ TeV}$$

For  $Y_B^{\text{obs}}$ ,  $T_I^\tau$  in the range 0.01 – 0.1

$$\Lambda/\sqrt{X_I^\tau} \lesssim 18 \text{ TeV} (0.01/T_I^\tau)^{1/2}$$

# Conclusions

- Baryon asymmetry can be produced with a tau CPV source.
- The CPV sources for the top and bottom cannot provide a large enough baryon asymmetry, due to EDM constraint.
- CPV source for the muon cannot provide large enough  $Y_B$  due to collider constraint  $h \rightarrow \mu\bar{\mu}$ .
- When multiple CPV sources (tau-t; tau-b) are present: cancellations to EDMs while enhancing  $Y_B > Y_B^{\text{obs}}$
- Smoking gun of this scenario is measuring CPV in Higgs boson decays to tau leptons.