

Contribution ID: 4

Type: Regular Talk (15'+5')

A low energy effective quark-gluon vertex to calculate the pseudoscalar meson spectrum

Monday 29 November 2021 16:25 (20 minutes)

Based on the generalized quantum electrodynamics expression for the Podolsky propagator, which preserves gauge invariance for massive photons, we propose a model for the massive gluon propagator that reproduces well-known features of established strong-interaction models in the framework of the Dyson-Schwinger equation. By adjusting the Podolsky mass and the coupling strength we thus construct a model with simple analytical properties known from perturbative theory, yet well suited to describe a confining interaction. We obtain solutions of the Dyson-Schwinger equation for the quark at spacelike momenta on the real axis as well as on the complex plane and solving the bound-state problem with the Bethe-Salpeter equation yields masses and weak decay constants of the π , K and η c in excellent agreement with experimental values, while the D and Ds are reasonably well described. The analytical simplicity of this effective interaction has the potential to be useful for phenomenological applications and may facilitate calculations in Minkowski space.

Authors: EL-BENNICH, Bruno (Universidade Cruzeiro do Sul); ROJAS, Eduardo (Universidad de Nariño); Dr RAMOS-ZAMBRANO, German (Universidad de Nariño.)

Presenter: ROJAS, Eduardo (Universidad de Nariño)

Session Classification: Heavy Flavour

Track Classification: Heavy Flavour