

NeMO-C 2024: Neutrinos y Materia Oscura en Colombia



Report of Contributions

Contribution ID: 1

Type: **not specified**

Dark matter stability and neutrinos nature from broken non-abelian discrete symmetries

Se estudia el rompimiento espontáneo de simetrías discretas no abelianas que dejen simetrías residuales conservadas. Con esto, podremos construir extensiones del modelo estándar que incluyan un sector oscuro que proporciona un candidato a materia oscura tipo WIMP. Al mismo tiempo que generan términos de masa para los neutrinos. Exploramos el espacio de parámetros para verificar la viabilidad del modelo y definir nuevos fenómenos observables detectables experimentalmente en el futuro cercano. Esto podría incluir procesos de violación de sabor en la desintegración del leptón tau y el quark top, los cuales están siendo analizados actualmente por el experimento CMS, así como la detección de posibles candidatos a materia oscura: directamente a través de las colaboraciones como DARWIN, e indirectamente mediante datos recopilados por CTA entre otros.

Authors: LAYANA, Andres (UCN); Dr BONILLA DÍAZ, Cesar (Universidad Católica del Norte)

Presenter: LAYANA, Andres (UCN)

Track Classification: Dark Matter

Contribution ID: 2

Type: **not specified**

Dark matter-electron interactions in a $L_{\mu} - L_{\tau}$ symmetry model

Wednesday 24 July 2024 10:30 (25 minutes)

We consider an $L_{\mu} - L_{\tau}$ abelian symmmetry extension of the Standard Model to derive spin-independent and spin-dependent interactions of fermion dark matter and electrons through the new gauge boson. We explore prospects with XLZD and OSCURA experiments to close the constraints in the parameter space able to explain simultaneously the recent measurement on the anomalous magnetic moment of the muon and the observed relic density of dark matter.

Author: OCHOA PEREZ, Fredy Alexander

Co-authors: HERRERA, Gonzalo (TUM, MPP); FIGUEROA, Pablo (Max Planck Institute for Physics)

Presenter: OCHOA PEREZ, Fredy Alexander

Track Classification: Dark Matter

Contribution ID: 3

Type: **not specified**

Thermal Dark Matter with Low-Temperature Reheating

Wednesday 24 July 2024 11:00 (25 minutes)

We explore the production of thermal dark matter (DM) candidates (WIMPs, SIMPs, ELDERs and Cannibals) during cosmic reheating. Assuming a general parametrization for the scaling of the inflaton energy density and the standard model (SM) temperature, we study the requirements for kinetic and chemical DM freeze-out in a model-independent way. For each of the mechanisms, up to two solutions that fit the entire observed DM relic density exist, for a given reheating scenario and DM mass. As an example, we assume a simple particle physics model in which DM interacts with itself and with SM through contact interactions. We find that low-temperature reheating can accommodate a wider range of couplings and larger masses than those permitted in the usual instantaneous high-temperature reheating. This results in DM solutions for WIMPs reaching masses as high as 10^{14} GeV, whereas for SIMPs and ELDERs, we can reach masses of 10^{13} GeV. Interestingly, current experimental data already constrain the enlarged parameter space of these models with low-reheating temperatures. Next-generation experiments could further probe these scenarios.

Authors: Dr DEKA, Kuldeep (New York University Abu Dhabi); LOSADA, Marta (NYUAD); BERNAL, Nicolás (New York University Abu Dhabi)

Presenter: BERNAL, Nicolás (New York University Abu Dhabi)

Track Classification: Dark Matter

Contribution ID: 4

Type: **not specified**

Effective dark matter

Wednesday 24 July 2024 09:30 (25 minutes)

Restrictions in Wilson coefficients for Dark matter simplified models

Author: Prof. RESTREPO QUINTERO, DIEGO ALEJANDRO

Presenter: Prof. RESTREPO QUINTERO, DIEGO ALEJANDRO

Track Classification: Dark Matter

Contribution ID: 5

Type: **not specified**

Collider analysis of the $U(1)_L$ globally gauged scotogenic model within compressed mass spectrum scenarios.

Thursday 25 July 2024 10:30 (25 minutes)

In this talk we present an variation of the Scotogenic Model, that extends the gauge group by a global $U(1)$ symmetry, and a singlet scalar that induces Spontaneous Symmetry Breaking to explain the origin of both Majorana Masses and Lepton number violation. Then, we make a brief analysis of the compatible parameter space for both fermionic and scalar dark matter, which can be considered in a compressed mass spectrum between the lightest fermionic and scalar states. Then, we make an analysis of the behavior of the production cross section of the DM candidate particle, for both Drell-Yan and Vector Boson Fusion mechanisms and different compressed mass spectra scenarios, as a function of the mass and compare it with the latest results given by the ATLAS and CMS experiments to establish the detection feasibility of the model at the LHC.

Author: FLÓREZ, Andrés (Universidad de Los Andes)

Presenter: FLÓREZ, Andrés (Universidad de Los Andes)

Track Classification: Neutrino Physics

Contribution ID: 6

Type: **not specified**

Phenomenology of a singlet-doublet-triplet scotogenic framework

Thursday 25 July 2024 11:00 (25 minutes)

I will present a triplet-scotogenic model, so called “T1-2G”, in order to provide a solution to neutrino masses, and dark matter, while satisfying the limit on lepton flavor violation processes. The viable region of the parameter space is determined through a Markov Chain Monte Carlo numerical code, and satisfies both dark matter observable and lepton flavor violation constraints.

Authors: HERRMANN, Bjorn; SARAZIN, Maud; DE NOYERS, Ugo (LAPTh / Université Savoie Mont Blanc)

Presenter: SARAZIN, Maud

Track Classification: Neutrino Physics

Contribution ID: 7

Type: **not specified**

Reopening the Z portal with semi-annihilations

Wednesday 24 July 2024 15:35 (25 minutes)

In one-component dark matter (DM) scenarios is commonly assumed that a scalar WIMP must either be part of an $SU(2)_L$ multiplet with zero hypercharge or have suppressed vector interactions with the Z gauge boson to circumvent stringent direct detection (DD) bounds. In this work, we demonstrate that multi-component scenarios with a dark scalar doublet exhibiting vector-like interactions with the Z boson are also compatible with bounds arising from DD searches. Specifically, we consider a simple extension of the Standard Model wherein the dark sector comprises a doublet and a complex singlet ϕ , both charged under a Z_6 symmetry. We find that semi-annihilation processes drastically reduce the relic abundance of the neutral component of the doublet, H^0 , sufficiently attenuating the effects of its large Z -mediated elastic scattering cross-section with nucleons to satisfy the DD constraints. Although the contribution of H^0 to the total relic abundance is nearly negligible, with ϕ dominating, both dark matter components are expected to be detectable in ongoing and future DD experiments. The viability of the model is tested against several theoretical and experimental constraints, resulting in a parameter space featuring a non-degenerate mass spectrum at the electroweak scale.

Author: Dr ZAPATA, Óscar (Universidad de Antioquia)

Co-authors: Ms DOMÍNGUEZ, María J. (Universidad de Antioquia); RODRIGUEZ CIFUENTES, OSCAR ANDRES

Presenter: RODRIGUEZ CIFUENTES, OSCAR ANDRES

Track Classification: Dark Matter

Contribution ID: 8

Type: **not specified**

Registration and Welcome

Wednesday 24 July 2024 08:00 (30 minutes)

Author: VANEGAS FORERO, David (Universidad de Medellin)

Presenter: VANEGAS FORERO, David (Universidad de Medellin)

Contribution ID: 9

Type: **not specified**

Dark Matter - Overview

Wednesday 24 July 2024 08:30 (55 minutes)

Author: BERNAL, Nicolás (New York University Abu Dhabi)

Presenter: BERNAL, Nicolás (New York University Abu Dhabi)

Contribution ID: **10**

Type: **not specified**

Neutrinos - Overview

Thursday 25 July 2024 08:30 (55 minutes)

Author: VANEGAS FORERO, David (Universidad de Medellin)

Presenter: VANEGAS FORERO, David (Universidad de Medellin)

Contribution ID: **11**

Type: **not specified**

Exercises

Thursday 25 July 2024 14:00 (55 minutes)

Author: BERNAL, Nicolás (New York University Abu Dhabi)

Presenter: BERNAL, Nicolás (New York University Abu Dhabi)

Contribution ID: 13

Type: **not specified**

$\mu - \tau$ reflection symmetry and magic neutrino mass matrix

Thursday 25 July 2024 11:30 (25 minutes)

The searches for an underlying pattern in neutrino masses have motivated different proposals for textures in the neutrino mass matrix, which is also related to particular arrangements of the mixing matrix. The current precise determinations of neutrino mixings have discarded some of the most studied proposals, such as the one involving a $\mu - \tau$ exchange symmetry. In this work, we investigate the relation of a still allowed $\mu - \tau$ reflection symmetry with the constraints of a magic pattern in the neutrino mass matrix. We show that both conditions cannot be fulfilled simultaneously in an exact but rather in an approximate way. These considerations may have some effects on the values of the CP-violating phases and in some observables like the neutrinoless double beta decay amplitude, which may be explored in future experiments.

Author: TOSTADO, Sergio (Universidad Santiago de Cali)**Co-author:** Dr RIVERA-AGUDELO, Diana C. (Universidad Santiago de Cali)**Presenter:** TOSTADO, Sergio (Universidad Santiago de Cali)**Track Classification:** Neutrino Physics

Contribution ID: 14

Type: **not specified**

Dark matter in QCD-like theories with a theta vacuum: cosmological and astrophysical implications

Wednesday 24 July 2024 15:05 (25 minutes)

QCD-like theories in which the dark matter (DM) of the Universe is hypothesized to be a thermal relic in the form of a dark pion has been extensively investigated, with most studies neglecting the CP-violating θ -angle associated with the topological vacuum. We point out that a non-vanishing θ could potentially trigger resonant number-changing processes giving rise to the observed relic density in agreement with perturbative unitarity as well as observations of clusters of galaxies. This constitutes a novel production mechanism of MeV DM and an alternative to those relying on the Wess-Zumino-Witten term. Moreover, for specific meson mass spectra, similar resonant scatterings serve as a realization of velocity-dependent self-interacting DM without a light mediator. Explicit benchmark models are presented together with a discussion of possible signals.

Author: ZAPATA, Oscar**Presenter:** ZAPATA, Oscar**Track Classification:** Dark Matter

Contribution ID: 15

Type: **not specified**

LArSoft and its applications in DUNE

Thursday 25 July 2024 09:30 (25 minutes)

In this talk we are going to provide an introduction to Fermilab computing and how to set up an account in Fermilab to use The Liquid Argon Software (LArSoft) and Its main functionalities. This tutorial will present the principal LArSoft modules used as tools in the simulation and reconstruction of different phenomena across liquid Argon time projection chambers, which are employed in neutrino experiments like DUNE. The main objective of this talk is to enable the audience to understand how a LArSoft simulation is conducted in DUNE, with special focus in the photon simulation module, while utilizing the ROOT framework for visualization results. Finally, we will present the broad context of data analysis tools used in the DUNE experiment.

Authors: MORENO, DEYWIS; TAMARA JARAMILLO, JOSE DAVID (Universidad Antonio Nariño)

Presenter: TAMARA JARAMILLO, JOSE DAVID (Universidad Antonio Nariño)

Track Classification: Neutrino Physics

Contribution ID: 16

Type: **not specified**

Breve introducción al sistema de detección de fotones del proyecto DUNE

Thursday 25 July 2024 16:30 (15 minutes)

El Proyecto DUNE cuyo objetivo es ahondar en el estudio y comprensión de los neutrinos consta de un generador de haz de neutrinos captados luego por un detector cercano y posteriormente por un detector lejano. Para detectar dichas partículas se hace necesario un gran volumen de material sensible y su correspondiente hardware orientado a amplificar y digitalizar la señal de los eventos físicos que caracterizan su interacción. En este orden de ideas, y acorde con los eventos físicos de interés, se hace necesario un sistema de detección de fotones. En esta exposición se hará una breve reseña del sistema de detección de fotones del detector lejano del proyecto DUNE en su detector de fase simple.

Author: OSSA SÁNCHEZ, Jorge Eduardo (Universidad de Medellín, Estudiante Doctorado en Modelación y Computación Científica)

Presenter: OSSA SÁNCHEZ, Jorge Eduardo (Universidad de Medellín, Estudiante Doctorado en Modelación y Computación Científica)

Track Classification: Neutrino Physics

Contribution ID: 17

Type: **not specified**

Fuzzy dark matter modeling of the dynamical behavior of nearby isolated dwarf galaxies.

Wednesday 24 July 2024 17:00 (25 minutes)

We investigate a model based on fuzzy dark matter (FDM) formalism to describe rotation curves of nearby isolated dwarf galaxies. Our study evaluates the consistency of the FDM behavior with high resolution rotation curves from the LITTLE THINGS 3D catalog, which contains dark matter dominated galaxies. We find that the phenomenological model incorporating a central soliton structure, as predicted by FDM, accurately represents the dynamics of the dwarf galaxies and their density profiles. Nevertheless, the halo mass function prediction of FDM is not compatible with the determination of the number of observable haloes in the distances of the nearby group of irregular dwarf galaxies. This discrepancy presents a catch-22 paradox for the minimal FDM model, particularly within the axion mass range that permits the formation of soliton cores.

Author: CASTILLO RAMIREZ, Andrés Fernando (IAC/ULL)**Presenter:** CASTILLO RAMIREZ, Andrés Fernando (IAC/ULL)**Track Classification:** Dark Matter

Contribution ID: 18

Type: **not specified**

Fenomenología de un modelo de masas de neutrinos de tipo dirac.

Thursday 25 July 2024 16:50 (15 minutes)

El modelo estándar de la física de partículas ha demostrado ser la teoría más exitosa de la historia, gracias a su impresionante poder predictivo y su capacidad para explicar una amplia gama de fenómenos naturales. No obstante, esta teoría no es definitiva y deja varias preguntas sin resolver, siendo una de las más intrigantes: ¿cómo adquieren masa los neutrinos? En este trabajo, abordamos esta cuestión proponiendo un modelo de masas de neutrinos de tipo Dirac en el contexto de la física más allá del modelo estándar. Presentamos el lagrangiano asociado a este modelo y exploramos el espacio de parámetros relevante. Posteriormente, derivamos la expresión para la matriz de masa de los neutrinos (m_ν) . Adicionalmente, encontramos una parametrización para sus acoplos, lo que permite restringir la nueva física usando resultados de física de neutrinos. Este modelo puede también ser explorado por otros decaimientos $h \rightarrow \gamma\gamma$ y $\mu \rightarrow e\gamma$, destacando sus implicaciones fenomenológicas y potenciales señales experimentales.

Author: MANCO, Sebastián (EIA University)**Co-authors:** BETANCUR RODRIGUEZ, Amalia (EIA University (CO)); PALACIO, Guillermo**Presenter:** MANCO, Sebastián (EIA University)**Track Classification:** Neutrino Physics

Contribution ID: 19

Type: **not specified**

Massive Kalb-Ramond field as Feebly-interacting dark matter

Wednesday 24 July 2024 16:05 (25 minutes)

The Kalb-Ramond (KR) field is an antisymmetric rank two tensor that appears in the context of string theory. In this work, we consider the massive KR field as a dark matter candidate and we study its production via the FIMP mechanism through the Higgs portal. We find that the KR field gives account for the relic density of dark matter today for a wide range of masses.

Presenter: Dr LONGAS, Robinson (UdeA)