

Magnetic field-dependence of the neutral pion mass in the linear sigma model coupled to quarks: The weak field case

Monday 6 May 2019 14:50 (20 minutes)

We compute the neutral pion mass dependence on a magnetic field in the weak field approximation at one-loop order. The calculation is carried out within the linear sigma model coupled to quarks and using Schwinger's proper-time representation for the charged particle propagators. We find that the neutral pion mass decreases with the field strength provided the boson self-coupling magnetic field corrections are also included. The calculation should be regarded as the setting of the trend for the neutral pion mass as the magnetic field is turned on.

Author: MANREZA PARET, Daryel (Facultad de Física, Universidad de La Habana)

Co-authors: Mr AYALA, Alejandro (ICN/UNAM); Mr L. S. FARIAS, Ricardo (Universidade Federal de Santa Maria, Santa Maria, RS 97105-900, Brazil.); Mr HERNANDEZ-ORTIZ, Saul (ICN/UNAM); Mr HERNANDEZ, Luis (ICN/UNAM); Mr ZAMORA, Renato (Centro de Investigacion y Desarrollo en Ciencias Aeroespaciales (CIDCA), Fuerza Aerea de Chile, Santiago, Chile.)

Presenter: MANREZA PARET, Daryel (Facultad de Física, Universidad de La Habana)

Track Classification: STARS