

## Kinematic Constraints on Spatial Curvature from Supernovae Ia and Cosmic Chronometers

This work has an interesting approach to estimate the spatial curvature  $\Omega_k$  from data independently of dynamical models is suggested, it was done through three kinematic parameterizations of the comoving distance ( $D_C(z)$ ) with second degree polynomial, of the Hubble parameter ( $H(z)$ ) with a second degree polynomial and of the deceleration parameter ( $q(z)$ ) with first order polynomial. All these parameterizations are done as function of redshift  $z$ . We used SNe Ia dataset from Pantheon compilation with 1048 distance moduli estimated on the range  $0.01 < z < 2.3$  with systematic and statistical errors and a compilation of 31  $H(z)$  data estimated from cosmic chronometers. The spatial curvature found for  $D_C(z)$  parametrization was  $\Omega_k = -0.49^{+0.14+0.29}_{-0.14-0.27}$ . The parametrization for deceleration parameter  $q(z)$  resulted in  $\Omega_k = -0.08^{+0.21+0.54}_{-0.27-0.45}$ . The  $H(z)$  parametrization had incompatibilities between  $H(z)$  and SNe Ia data, so these analyses were not combined. The  $q(z)$  parametrization is compatible with the spatially flat Universe as predicted by many inflation models and data from CMB, while the  $D_C(z)$  parametrization favored a slightly closed Universe. This type of analysis may be interesting as it avoids any bias because it does not depend on assumptions about the matter content for estimating  $\Omega_k$ .

**Authors:** VALENTIM, Rodolfo (UNIFESP - Diadema); Prof. JESUS, José Fernando (Universidade Estadual Paulista - UNESP); MALHEIRO, Manuel (ITA); Dr MORAES, P. H.R.S (IAG - USP)

**Presenter:** VALENTIM, Rodolfo (UNIFESP - Diadema)

**Session Classification:** DM, DE, GWs, BHs, GRAVITATION, GALAXIES, QCD, LIFE, GRBs, COSMOLOGY, OA, KT