



Contribution ID: 436

Type: **Poster**

A curved jet model for the synchrotron emission of the BL Lac object PG 1553+113.

Tuesday 8 December 2015 18:39 (3 minutes)

We report on the results of a multifrequency campaign on the BL Lac object PG 1553+113 that was organized by the Whole Earth Blazar Telescope (WEBT) Collaboration in 2013 April-August. Nineteen optical, two near-IR, and three radio telescopes monitored the source to follow its behaviour at low energies during and around the high-energy observations by the MAGIC telescopes in April-July. A general bluer-when-brighter trend characterizes the optical emission.

We also analyse the UV and X-ray data acquired by the Swift and XMM-Newton satellites in the same period and compare them with previous observations.

The long XMM-Newton exposure reveals a curved X-ray spectrum, which shows a harder-when-brighter behaviour on long time scales.

In the spectral energy distribution (SED), the XMM-Newton near-UV spectrum is hard, while Swift data display a softer shape that is confirmed by previous HST-COS and IUE observations.

An estimate of the synchrotron peak through polynomial fits to the optical-X-ray SED suggests that it lies in the 4-30 eV energy range, with a general increase with X-ray brightness.

However, the UV and X-ray spectra do not connect smoothly. We propose an interpretation of the SED shape and variability in terms of orientation changes of the helical structure of an inhomogeneous jet.

Authors: Dr STAMERRA, Antonio (INAF-Osservatorio Astrofisico di Torino, Italy); Dr RAITERI, Claudia M. (INAF-Osservatorio Astrofisico di Torino, Italy); Dr VILLATA, Massimo (INAF-Osservatorio Astrofisico di Torino)

Presenter: Dr STAMERRA, Antonio (INAF-Osservatorio Astrofisico di Torino, Italy)

Session Classification: 14 - Disks and jets