28th Texas Symposium on Relativistic Astrophysics



Contribution ID: 433

Type: Talk

First numerical simulations of the chiral MHD dynamo effect

Monday 7 December 2015 14:42 (21 minutes)

Under extreme conditions, e.g. at high temperatures like in the early Universe, the usual magnetohydrodynamical (MHD) equations need to be extended. The origin of the modification is the asymmetry of the chemical potential of right- and left-handed fermions. To describe the evolution of a plasma, additional terms as well as new equations for the chiral chemical potential have to be included. We have implemented these extensions in the Pencil Code, which is a high-order finite-difference code developed for solving compressible MHD. We study laminar dynamos and find an exponential increase of the magnetic energy with the growth rate depending on the chemical potential. We further analyze the evolution of magnetic fields in the presence of turbulence. Potentially, chirality has interesting effects on the properties of magnetic fields in the early Universe and in particular on the evolution of magnetic helicity which is directly coupled to the chiral chemical potential.

Author: SCHOBER, Jennifer

Co-authors: BOYARSKY, Alexey (CERN); BRANDENBURG, Axel (Nordita); ROGACHEVSKII, Igor (Ben Gurion University of the Negev); RUCHAYSKIY, Oleg (Ecole Polytechnique Federale de Lausanne (CH))

Presenter: SCHOBER, Jennifer

Session Classification: 10 - Cosmic magnetic fields