



Contribution ID: 181

Type: Talk

The role of general relativity and reconnection in pulsar radiation

Sunday 6 December 2015 14:42 (20 minutes)

Pulsars shine throughout the electromagnetic spectrum, from radio waves to energetic gamma rays. The radio emission is thought to originate from the discharge of the polar-cap and the formation of copious electron-positron pairs. Gamma rays are traditionally associated with particle acceleration in electrostatic gaps within the light cylinder. The recent development of global Particle-In-Cell (PIC) simulations of the pulsar magnetosphere enables to test these scenarios self-consistently. We show that general relativistic effects, most notably frame-dragging, are essential to ignite pair creation in the polar cap for low-inclination pulsars, and hence enable pulsars to emit radio waves. In addition, three-dimensional radiative PIC simulations indicate that the current sheet that forms beyond the light cylinder is the main site of particle acceleration in plasma-filled pulsars, instead of gaps within the co-rotating magnetosphere. Relativistic reconnection dissipates the magnetic energy which is then converted to energetic particles and high-energy synchrotron radiation. We present self-consistent modeling of pulsar gamma-ray lightcurves and spectra obtained directly from the kinetic simulations, and discuss the results in the context of observed gamma-ray pulsars.

Author: CERUTTI, Benoit (Princeton University)

Co-authors: PHILIPPOV, Alexander (Princeton University); SPITKOVSKY, Anatoly (Princeton University)

Presenter: CERUTTI, Benoit (Princeton University)

Session Classification: 18 - Gal. accel. & pulsars