



Contribution ID: 44

Type: **Poster**

Effects of the second-order vector mode on weak lensing signals

Tuesday 8 December 2015 17:41 (3 minutes)

Vector mode of cosmological perturbation theory imprints characteristic signals on the weak lensing signals such as curl- and B-modes which are never imprinted by the scalar mode.

However, the vector mode is neglected in the standard first-order cosmological perturbation theory since it only has a decaying mode.

This situation changes if the cosmological perturbation theory is expanded up to second order.

The second-order vector and tensor modes are inevitably induced by the product of the first-order scalar modes.

We study the effect of the second-order vector mode on the weak lensing curl- and B-modes.

We find that the curl-mode induced by the second-order vector mode is comparable to that induced by the primordial gravitational waves with the tensor-to-scalar ratio $r = 0.1$ at $\ell \approx 200$.

In this case, the curl-mode induced by the second-order vector mode dominates at $\ell > 200$.

Furthermore, the B-mode cosmic shear induced by the second-order vector mode dominates on almost all scales.

However, we find that the observational signatures of the second-order vector and tensor modes cannot exceed the expected noise

of ongoing and upcoming weak lensing measurements.

We conclude that the curl- and B-modes induced by the second-order vector and tensor modes are unlikely to be detected in future experiments.

Author: Mr SAGA, Shohei (Naogya university)

Co-authors: Dr YAMAUCHI, Daisuke (The University of Tokyo); Dr ICHIKI, Kiyotomo (Nagoya university)

Presenter: Mr SAGA, Shohei (Naogya university)

Session Classification: 12 - Gravitational lensing