
Signal and background in FLArE

Jianming Bian, Wenjie Wu

University of California, Irvine



June 9, 2022



UCIRVINE


Detector configuration in Geant4

| | LArTPC | HadCal | MuonFinder |
|-------------|----------|-------------|-------------|
| Length (mm) | 0 - 7000 | 7250 - 8300 | 8300 - 9340 |



Simulation setup

- The vertices of neutrino interactions are uniformly distributed in the FV region (1x1x7 m)
- No angular smearing for the neutrino beam, all pointed at +z direction
- Same amount of neutrino interactions were simulated for ν_e , ν_μ , and ν_τ
 - flux ratio is $\sim 100:400:1$



Signal and background

- Only consider beam neutrino background for now
- Decay modes of the tau lepton
 - τ_e : taus decay to electrons
 - τ_μ : taus decay to muons
 - τ_{had} : taus decay to hadrons
- Major background of τ_μ signal: ν_μ CC events
- Major background of τ_e signal: ν_e CC events
- Major background of τ_{had} signal: NC scattering events from all neutrinos

TABLE I. Dominant decay modes of τ^- . All decays involving kaons, as well as other subdominant decays, are in the “Other” category.

| Decay mode | Branching ratio |
|------------------------------------|-----------------|
| Leptonic | 35.2% |
| $e^- \bar{\nu}_e \nu_\tau$ | 17.8% |
| $\mu^- \bar{\nu}_\mu \nu_\tau$ | 17.4% |
| Hadronic | 64.8% |
| $\pi^- \pi^0 \nu_\tau$ | 25.5% |
| $\pi^- \nu_\tau$ | 10.8% |
| $\pi^- \pi^0 \pi^0 \nu_\tau$ | 9.3% |
| $\pi^- \pi^- \pi^+ \nu_\tau$ | 9.0% |
| $\pi^- \pi^- \pi^+ \pi^0 \nu_\tau$ | 4.5% |
| Other | 5.7% |








FIG. 2. Pictorial representation of hadronic tau (upper left) and leptonic tau (upper right) signals, and their corresponding backgrounds (lower).

[10.1103/PhysRevD.102.053010](https://arxiv.org/abs/10.1103/PhysRevD.102.053010)




Signatures of the final state particles

- Stable particles in the final state, including particles from tau decay
- τ_{had} have at least $1\pi^-$ in the final state
 - τ_{had} have more (π^-, π^0) in the final state
- Neutrinos in the final state are invisible to the detector, contributing to the missing energy
 - Almost all $\nu_{\mu} \text{CC}, \nu_e \text{CC}$ have zero neutrino in the final state
 - NC events and τ_{had} have 1 neutrino, τ_{μ} and τ_e have 2 neutrinos






Deposited energy of the longest prong

- τ_μ and ν_μ CC are distinctive from other channels





Some other features

- τ_μ and ν_μ CC have relatively longer shower length
- Averaged $\langle dE/dx \rangle$ can also be used to select τ_μ and ν_μ CC




Missing transverse momentum

- τ_μ have more neutrinos in the final state than ν_μ CC, thus more missing momentum in the transverse plane



Energy of the leading π^-

- τ_{had} has larger branch ratio than τ_{μ} and τ_e , there is potential to be a good channel to select ν_{τ}
- τ_{had} generally have a more energetic π^- in the final state





Next steps

- Some of the features are promising to differentiate ν_τ CC from other
 - $\langle dE/dx \rangle$, Missing momentum, leading π^- energy, ...
- Trying to look for more features
 - π^0 s in τ_{had} decay
 - ...
- These features could be used to do event selections, and make efficiency estimation, et.al.
 - A BDT for event identification could be a good start

Backup

Detector configuration in Geant4

| | LArTPC | HadCal | MuonFinder |
|-------------|----------|-------------|-------------|
| Length (mm) | 0 - 7000 | 7250 - 8300 | 8300 - 9340 |





LArTPC

HadCal

MuonFinder




Work in progress

- Save all the hit information from G4 simulation
 - As the energy is very high, there is a large amount of hits for each event (~TB for 10000 neutrino events)
- Will do more analysis on the new MC data
 - Study the feature of all stable final state particles from the neutrino interaction
 - Event classification, background rejection




ν_τ s in the detector

- Neutrino vertices are uniformly distributed in a 1x1x7 meter volume
- Neutrino energy/Interaction mode/FSL come from GENIE v3_00_06k
 - Flux comes from *Weidong Bai, et. al. 2112.11605*



τ^- s in the detector






| Decay mode | Branching ratio |
|------------------------------------|-----------------|
| Leptonic | 35.2% |
| $e^- \bar{\nu}_e \nu_\tau$ | 17.8% |
| $\mu^- \bar{\nu}_\mu \nu_\tau$ | 17.4% |
| Hadronic | 64.8% |
| $\pi^- \pi^0 \nu_\tau$ | 25.5% |
| $\pi^- \nu_\tau$ | 10.8% |
| $\pi^- \pi^0 \pi^0 \nu_\tau$ | 9.3% |
| $\pi^- \pi^- \pi^+ \nu_\tau$ | 9.0% |
| $\pi^- \pi^- \pi^+ \pi^0 \nu_\tau$ | 4.5% |
| other | 5.7% |

<https://arxiv.org/pdf/2007.00015.pdf>




ν_μ s in the detector

- Neutrino vertices are uniformly distributed in a 1x1x7 meter volume
- Neutrino energy/Interaction mode/FSL come from GENIE v3_00_06k
 - Flux comes from *Felix Kling, et. al. 2105.08270*




ν_e s in the detector

- Neutrino vertices are uniformly distributed in a 1x1x7 meter volume
- Neutrino energy/Interaction mode/FSL come from GENIE v3_00_06k
 - Flux comes from *Felix Kling, et. al. 2105.08270*




Neutrino flux



Felix Kling, et. al. [2105.08270](#)
[Github](#)

FLArE10, 620m downstream from IP, 3000/fb


x Luminosity / 2



Weidong Bai, et. al. [2112.11605](#)
 Figure 12, Table 5

eta > 6.9 (radius 1 m at a distance of 480 m from IP)

GENIE simulation: muon spectrum



$\nu_\mu \rightarrow \mu^-$
Mean: 343.1 GeV
RMS: 377.6 GeV


$\nu_\tau \rightarrow \tau^- \rightarrow \mu^-$
Mean: 146.0 GeV
RMS: 201.0 GeV

Felix Kling, et. al. [2105.08270](#)

Muon energy spectrum, area normalized

Muon from tau decay is softer

GENIE simulation: muon spectrum



Bai, $\nu_\tau \rightarrow \tau^- \rightarrow \mu^-$
Mean: 102.9 GeV
RMS: 136.7 GeV

Kling, $\nu_\tau \rightarrow \tau^- \rightarrow \mu^-$
Mean: 146.0 GeV
RMS: 201.0 GeV

Muon energy spectrum, area normalized