

# Approximate alignment without decoupling in the 2HDM naturally

7th RISE Collaboration  
Workshop in Helsinki

Howard E. Haber  
27 May 2019



Congratulations to our Finnish hosts  
for a well-timed conference!



# Outline

- Extended Higgs sector—motivations and constraints
- Achieving a SM-like Higgs boson in the 2HDM
- Approximate Higgs alignment via an approximate symmetry
- Possible symmetries of the 2HDM scalar potential
- Extending the symmetry to the Yukawa sector via mirror fermions
- Natural Higgs alignment without decoupling in the 2HDM
- Future work

This work is based on P. Draper, A. Ekstedt and H.E. Haber, in preparation.  
The central idea originated in P. Draper, H.E. Haber and J.T. Ruderman, JHEP **1606**, 124 (2016) [arXiv:1605.03237].

## Why not an extended Higgs sector?

- The fermion and gauge boson sectors of the Standard Model (SM) are not of minimal form (“Who ordered that?”). So, why should the spin-0 (scalar) sector be minimal?
- Extended Higgs sectors can provide a dark matter candidate.
- Extended Higgs sectors can modify the electroweak phase transition and facilitate baryogenesis.
- Extended Higgs sectors can enhance vacuum stability.
- Models of new physics beyond the SM often require additional scalar Higgs states. E.g., two Higgs doublets are required in the minimal supersymmetric extension of the SM (MSSM).

## Extended Higgs sectors are highly constrained

- The electroweak  $\rho$  parameter is very close to 1.
- One neutral Higgs scalar of the extended Higgs sector must be SM-like (and identified with the Higgs boson at 125 GeV).
- At present, only one Higgs scalar has been observed.
- Higgs-mediated flavor-changing neutral currents (FCNCs) are suppressed.
- Charged Higgs exchange at tree level (e.g. in  $\bar{B} \rightarrow D^{(*)} \tau^- \bar{\nu}_\tau$ ) and at one-loop (e.g. in  $b \rightarrow s \gamma$ ) can significantly constrain the charged Higgs mass and the Yukawa couplings.
- If the scale that governs the non-SM-like Higgs bosons is close to the electroweak scale, is the naturalness problem of electroweak symmetry breaking exacerbated?

## A SM-like Higgs boson in an extended Higgs sector

Let us focus on the two-Higgs doublet model (2HDM) as a prototype for an extended Higgs sector. Consider the 2HDM scalar potential,

$$\begin{aligned} \mathcal{V} = & m_{11}^2 \Phi_1^\dagger \Phi_1 + m_{22}^2 \Phi_2^\dagger \Phi_2 - [m_{12}^2 \Phi_1^\dagger \Phi_2 + \text{h.c.}] + \frac{1}{2} \lambda_1 (\Phi_1^\dagger \Phi_1)^2 \\ & + \frac{1}{2} \lambda_2 (\Phi_2^\dagger \Phi_2)^2 + \lambda_3 (\Phi_1^\dagger \Phi_1) (\Phi_2^\dagger \Phi_2) + \lambda_4 (\Phi_1^\dagger \Phi_2) (\Phi_2^\dagger \Phi_1) \\ & + \left\{ \frac{1}{2} \lambda_5 (\Phi_1^\dagger \Phi_2)^2 + [\lambda_6 (\Phi_1^\dagger \Phi_1) + \lambda_7 (\Phi_2^\dagger \Phi_2)] \Phi_1^\dagger \Phi_2 + \text{h.c.} \right\}. \end{aligned}$$

The  $\Phi_i$  are hypercharge  $Y = 1$  doublets. After minimizing the scalar potential,  $\langle \Phi_i^0 \rangle = v_i / \sqrt{2}$  (for  $i = 1, 2$ ) with  $v \equiv (|v_1|^2 + |v_2|^2)^{1/2} = 2m_W/g = 246$  GeV.

Define the scalar doublet fields of the **Higgs basis**,

$$H_1 = \begin{pmatrix} H_1^+ \\ H_1^0 \end{pmatrix} \equiv \frac{v_1^* \Phi_1 + v_2^* \Phi_2}{v}, \quad H_2 = \begin{pmatrix} H_2^+ \\ H_2^0 \end{pmatrix} \equiv \frac{-v_2 \Phi_1 + v_1 \Phi_2}{v},$$

such that  $\langle H_1^0 \rangle = v/\sqrt{2}$  and  $\langle H_2^0 \rangle = 0$ . The Higgs basis is uniquely defined up to an overall rephasing,  $H_2 \rightarrow e^{i\chi} H_2$ .

## The Higgs basis and the alignment limit

The neutral scalar  $H_1^0$  is *aligned* in field space with the vacuum expectation value  $v$ . If  $\sqrt{2} H_1^0 - v$  were a mass eigenstate, then its tree-level properties would coincide with the Higgs boson of the SM.

In the Higgs basis, the scalar potential is given by:

$$\begin{aligned} \mathcal{V} = & Y_1 H_1^\dagger H_1 + Y_2 H_2^\dagger H_2 + [Y_3 H_1^\dagger H_2 + \text{h.c.}] + \frac{1}{2} Z_1 (H_1^\dagger H_1)^2 \\ & + \frac{1}{2} Z_2 (H_2^\dagger H_2)^2 + Z_3 (H_1^\dagger H_1) (H_2^\dagger H_2) + Z_4 (H_1^\dagger H_2) (H_2^\dagger H_1) \\ & + \left\{ \frac{1}{2} Z_5 (H_1^\dagger H_2)^2 + [Z_6 (H_1^\dagger H_1) + Z_7 (H_2^\dagger H_2)] H_1^\dagger H_2 + \text{h.c.} \right\}, \end{aligned}$$

After minimizing the scalar potential,  $Y_1 = -\frac{1}{2} Z_1 v^2$  and  $Y_3 = -\frac{1}{2} Z_6 v^2$ .

Remark:

Exact alignment corresponds to  $Z_6 = 0$ , which implies no  $H_1^0 - H_2^0$  mixing.

For simplicity, assume a CP-conserving scalar potential (where all Higgs basis parameters can be chosen real). The CP-even Higgs squared-mass matrix is,

$$\mathcal{M}_H^2 = \begin{pmatrix} Z_1 v^2 & Z_6 v^2 \\ Z_6 v^2 & m_A^2 + Z_5 v^2 \end{pmatrix}.$$

where  $m_A$  is the mass of the CP-odd Higgs scalar.

The CP-even Higgs bosons are  $h$  and  $H$  with  $m_h \leq m_H$ . Approximate alignment arises two limiting cases:

1.  $m_A^2 \gg (Z_1 - Z_5)v^2$ . This is the *decoupling limit*, where  $h$  is SM-like and  $m_A^2 \sim m_H^2 \sim m_{H^\pm}^2 \gg m_h^2 \simeq Z_1 v^2$ .
2.  $|Z_6| \ll 1$ . Then,  $h$  is SM-like if  $m_A^2 + (Z_5 - Z_1)v^2 > 0$ . Otherwise,  $H$  is SM-like. This is alignment with or without decoupling, depending on the value of  $m_A$ . The boundary between these two regions is fuzzy.

In particular, the CP-even neutral scalar mass eigenstates are:



$$\begin{pmatrix} H \\ h \end{pmatrix} = \begin{pmatrix} c_{\beta-\alpha} & -s_{\beta-\alpha} \\ s_{\beta-\alpha} & c_{\beta-\alpha} \end{pmatrix} \begin{pmatrix} \sqrt{2} \operatorname{Re} H_1^0 - v \\ \sqrt{2} \operatorname{Re} H_2^0 \end{pmatrix},$$

where  $c_{\beta-\alpha} \equiv \cos(\beta - \alpha)$  and  $s_{\beta-\alpha} \equiv \sin(\beta - \alpha)$  are defined in terms of the mixing angle  $\alpha$  that diagonalizes the CP-even Higgs squared-mass matrix when expressed in the  $\Phi_1$ - $\Phi_2$  basis of scalar fields,  $\{\sqrt{2} \operatorname{Re} \Phi_1^0 - v_1, \sqrt{2} \operatorname{Re} \Phi_2^0 - v_2\}$ , and  $\tan \beta \equiv v_2/v_1$ .

Since the SM-like Higgs boson must be approximately  $\sqrt{2} \operatorname{Re} H_1^0 - v$ , it follows that

- $h$  is SM-like if  $|c_{\beta-\alpha}| \ll 1$  (alignment with or without decoupling, depending on the value of  $m_A$ ),
- $H$  is SM-like if  $|s_{\beta-\alpha}| \ll 1$  (alignment without decoupling).

# LHC constraints on alignment in the 2HDM



Taken from ATLAS-CONF-2019-005 (March 20, 2019), under the assumption that  $h(125)$  is the lighter of the two CP-even scalars.

## Achieving a SM-like Higgs boson in the 2HDM

- In the decoupling limit,  $m_h \ll m_H, m_A, m_{H^\pm}$ . The SM is the effective low energy theory below the mass scale of the Higgs basis field  $H_2$ , and  $h$  is the SM-like Higgs boson.
- **The inert doublet model (IDM)**: There is a  $\mathbb{Z}_2$  symmetry in the Higgs basis such that  $H_2 \rightarrow -H_2$  is the only  $\mathbb{Z}_2$ -odd field. Then  $Z_6 = 0$ , and the tree-level properties of  $\sqrt{2}H_1 - v$  coincide with the SM Higgs boson. That is, tree-level alignment is exact. Deviations from SM behavior can appear at loop level due to the virtual exchange of the scalar states that reside in  $H_2$ . The lightest of the  $\mathbb{Z}_2$ -odd scalars is a dark matter candidate.
- Approximate alignment without decoupling. If present,
  - is this a result of an accidental choice of model parameters?
  - is this a consequence of an approximate (softly-broken) symmetry?  
(The latter is not possible in the IDM.)

## Possible symmetries of the 2HDM scalar potential

A complete classification of possible Higgs family and generalized CP symmetries of the scalar potential (in the  $\Phi_1$ – $\Phi_2$  basis) has been obtained.<sup>1</sup>

symmetry	$m_{22}^2$	$m_{12}^2$	$\lambda_2$	$\lambda_3$	$\lambda_4$	$\lambda_5$	$\lambda_6$	$\lambda_7$
$\mathbb{Z}_2$		0					0	0
$\Pi_2$	$m_{11}^2$	real	$\lambda_1$			real		$\lambda_6^*$
U(1)		0				0	0	0
SO(3)	$m_{11}^2$	0	$\lambda_1$		$\lambda_1 - \lambda_3$	0	0	0
CP		real				real	real	real
GCP2	$m_{11}^2$	0	$\lambda_1$					$-\lambda_6$
GCP3	$m_{11}^2$	0	$\lambda_1$			$\lambda_1 - \lambda_3 - \lambda_4$ (real)	0	0

### Remarks:

1.  $\Pi_2$  symmetry is equivalent to  $\mathbb{Z}_2$  symmetry in a different basis.
2. Simultaneous  $\mathbb{Z}_2$  and  $\Pi_2 \iff$  GCP2 in a different basis.
3. Simultaneous  $U(1)_{PQ}$  and  $\Pi_2 \iff$  GCP3 in a different basis.

---

<sup>1</sup>I.P. Ivanov, Phys. Rev. D **77**, 015017 (2008) [arXiv:0710.3490]; P.M. Ferreira, H.E. Haber and J.P. Silva, Phys. Rev. D **79**, 116004 (2009) [arXiv:0902.1537].

## A symmetry origin for alignment without decoupling

Consider the CP-conserving 2HDM. The scalar potential parameters in the  $\Phi_1$ – $\Phi_2$  basis are related to the corresponding Higgs basis parameters; e.g.,

$$Y_3 = \frac{1}{2}(m_{22}^2 - m_{11}^2)s_{2\beta} - m_{12}^2c_{2\beta}.$$

If  $m_{11}^2 = m_{22}^2$  and  $m_{12}^2 = 0$ , then  $Y_3 = 0$ . The scalar potential minimum condition ( $Y_3 = -\frac{1}{2}Z_6v^2$ ) then yields  $Z_6 = 0$ , i.e. exact alignment.<sup>2</sup> This leads to three possible symmetry choices:

symmetry	$m_{22}^2$	$m_{12}^2$	$\lambda_2$	$\lambda_3$	$\lambda_4$	$\lambda_5$	$\lambda_6$	$\lambda_7$
GCP2	$m_{11}^2$	0	$\lambda_1$					$-\lambda_6$
GCP3	$m_{11}^2$	0	$\lambda_1$			$\lambda_1 - \lambda_3 - \lambda_4$ (real)	0	0
SO(3)	$m_{11}^2$	0	$\lambda_1$		$\lambda_1 - \lambda_3$	0	0	0

Unfortunately, none of these symmetries can be extended to the Yukawa interactions without generating a massless quark or some other phenomenologically untenable feature.<sup>3</sup>

<sup>2</sup>See, e.g., P.S. Bhupal Dev and A. Pilaftsis, JHEP **1412**, 024 (2014) [Erratum: JHEP **1511**, 147 (2015)].

<sup>3</sup>P.M. Ferreira and J.P. Silva, Eur. Phys. J. C **69**, 45 (2010).

## The GCP-symmetric 2HDM with mirror fermions

The 2HDM with a GCP2 [GCP3]-symmetric scalar potential can be realized in another basis as a  $\mathbb{Z}_2 \otimes \Pi_2$  [ $U(1)_{PQ} \otimes \Pi_2$ ] discrete symmetry, where

$$m_{11}^2 = m_{22}^2, \quad \lambda_1 = \lambda_2, \quad \lambda_5 \text{ real } [\lambda_5 = 0], \quad m_{12}^2 = \lambda_6 = \lambda_7 = 0.$$

To extend this symmetry to the Yukawa sector, we introduce mirror fermions  $U$  and  $\bar{U}$ . SM two-component fermions are denoted by lower case letters (e.g. doublet fields  $q = (u, d)$  with  $Y = 1/3$  and singlet fields  $\bar{u}$  with  $Y = -4/3$ ); mirror singlet two-component fermions by upper case letters. Note that  $Y_{\bar{u}} = Y_{\bar{U}} = -Y_U$ . Under the symmetries,<sup>4</sup>

symmetry	$\Phi_1$	$\Phi_2$	$q$	$\bar{u}$	$\bar{U}$	$U$
$\mathbb{Z}_2$	$\Phi_1$	$-\Phi_2$	$q$	$-\bar{u}$	$\bar{U}$	$U$
$\Pi_2$	$\Phi_2$	$\Phi_1$	$q$	$\bar{U}$	$\bar{u}$	$U$
$U(1)$	$e^{-i\theta}\Phi_1$	$e^{i\theta}\Phi_2$	$q$	$e^{-i\theta}\bar{u}$	$e^{i\theta}\bar{U}$	$e^{-i\theta}U$

<sup>4</sup>The down-type fermions and leptons can also be included by introducing the appropriate mirror fermions.

The Yukawa couplings consistent with the  $\mathbb{Z}_2 \otimes \Pi_2$  [ $U(1)_{PQ} \otimes \Pi_2$ ] symmetry and the  $SU(2) \times U(1)_Y$  gauge symmetry are

$$\mathcal{L}_{\text{Yuk}} \supset y_t (q\Phi_2 \bar{u} + q\Phi_1 \bar{U}) + \text{h.c.}$$

The model is not phenomenologically viable due to



- experimental limits on mirror fermion masses
- existence of a massless scalar if  $U(1)_{PQ}$  is spontaneously broken

Thus, we introduce  $SU(2) \times U(1)_Y$  preserving mass terms associated with mirror fermions,

$$\mathcal{L}_{\text{mass}} \supset M_U \bar{U}U + M_u \bar{u}U + \text{h.c.}$$

The  $\mathbb{Z}_2$  [ $U(1)_{PQ}$ ] symmetry is preserved by the  $\bar{U}U$  mass term, whereas it is explicitly broken by the  $\bar{u}U$  mass term. The  $\Pi_2$  discrete symmetry is also explicitly broken if  $M_U \neq M_u$ . In all cases the symmetry breaking is soft, so that corrections to the scalar potential parameters are protected from quadratic sensitivity to the cutoff scale  $\Lambda$  of the theory.

## Effects of the broken symmetries



$$\Delta m^2 \equiv m_{22}^2 - m_{11}^2 \sim \kappa(M_U^2 - M_u^2) - \frac{3y_t^2(M_U^2 - M_u^2)}{4\pi^2} \ln(\Lambda/M) ,$$


where  $M \equiv (M_U^2 + M_u^2)^{1/2}$ . The above result includes a finite threshold corrections proportional to  $\kappa$ . Note that when  $M_U = M_u$ , the  $\Pi_2$  discrete symmetry is unbroken and hence the relation  $m_{11}^2 = m_{22}^2$  is protected. Likewise,

$$m_{12}^2 \sim \kappa_{12}M_U M_u + \frac{3y_t^2 M_U M_u}{4\pi^2} \ln(\Lambda/M) ,$$

which includes a finite threshold corrections proportional to  $\kappa_{12}$ .

Integrating out the mirror fermions below the scale  $M$ , one generates a splitting between  $\lambda_1$  and  $\lambda_2$  and nonzero values of  $\lambda_{5,6,7}$ .

For example, above the scale  $M$ , the diagrams



contribute equally to  $\lambda_2(\Phi_2^\dagger\Phi_2)^2$  and  $\lambda_1(\Phi_1^\dagger\Phi_1)^2$ , respectively. Below the scale  $M$ , the diagrams with internal  $U$  lines decouple, which then yields

$$\Delta\lambda \equiv |\lambda_1 - \lambda_2| \sim \frac{3y_t^4}{4\pi^2} \log(M/m_t) \sim 0.1,$$

for  $M \sim \mathcal{O}(1 \text{ TeV})$ . This is a small correction, which in first approximation can be neglected in our analysis.

Likewise, explicit breaking of the  $\mathbb{Z}_2$   $[\text{U}(1)_{\text{PQ}}]$  symmetry will generate small nonzero values of  $[\lambda_5]$ ,  $\lambda_6$  and  $\lambda_7$ .

## Top quark–mirror quark mixing

After electroweak symmetry breaking, the fermion mass eigenstates are obtained by Takagi-diagonalization of the following  $4 \times 4$  mass matrix.

$$-\mathcal{L}_{\text{mass}} = \frac{1}{2}(u \quad U \quad \bar{u} \quad \bar{U}) \begin{pmatrix} 0 & 0 & m_2 & m_1 \\ 0 & 0 & M_u & M_U \\ m_2 & M_u & 0 & 0 \\ m_1 & M_U & 0 & 0 \end{pmatrix} \begin{pmatrix} u \\ U \\ \bar{u} \\ \bar{U} \end{pmatrix} + \text{h.c.},$$

where  $m_1 \equiv y_t v_1 / \sqrt{2}$  and  $m_2 \equiv y_t v_2 / \sqrt{2}$ . States with the same electric charge, i.e.  $\{u, U\}$  and  $\{\bar{u}, \bar{U}\}$ , can separately mix (with mixing angles  $\theta_L$  and  $\theta_R$ , respectively). This yields two Dirac fermions—the top quark  $t$  and its mirror  $T$ , with squared-masses,

$$\left\{ \begin{matrix} M_T^2 \\ m_t^2 \end{matrix} \right\} = \frac{1}{2} \left[ m^2 + M^2 \pm \sqrt{(m^2 + M^2)^2 - 4(m_1 M_u - m_2 M_U)^2} \right],$$

where  $m \equiv y_t v / \sqrt{2}$  and  $M^2 \equiv M_U^2 + M_u^2$ , and  $\tan \theta_L = (m_t / m_T) \tan \theta_R$ .

## The Higgs sector of the softly-broken GCP-symmetric 2HDM

---

The important parameters of the scalar potential are:

$$m^2 \equiv \frac{1}{2}(m_{11}^2 + m_{22}^2), \quad \Delta m^2 \equiv m_{22}^2 - m_{11}^2, \quad R \equiv \frac{\lambda_{345}}{\lambda}, \quad m_{12}^2,$$

where  $\lambda_{345} \equiv \lambda_3 + \lambda_4 + \lambda_5$ . We impose  $\lambda > 0$  and  $R > -1$  to ensure that the vacuum is bounded from below. Solving for the potential minimum yields,

$$2m^2 = \bar{m}^2 - \frac{1}{2}\lambda v^2(1 + R), \quad \Delta m^2 = \epsilon \left( \bar{m}^2 + \frac{1}{2}\lambda v^2(1 - R) \right),$$

where  $\bar{m}^2 \equiv 2m_{12}^2/\sin 2\beta$  and

$$\tan \beta \equiv \frac{v_2}{v_1} = \sqrt{\frac{1 - \epsilon}{1 + \epsilon}}, \quad \text{where } \epsilon \equiv \cos 2\beta.$$

The positivity of  $v_1^2$  and  $v_2^2$  requires  $|\epsilon| < 1$ .

## Approximate alignment without decoupling

The relevant Higgs basis parameters are given by,

$$\begin{aligned} Z_1 &= \frac{1}{2}\lambda[1 + R + \epsilon^2(1 - R)] , \\ m_A^2 + Z_5 v^2 &= \bar{m}^2 + \frac{1}{2}\lambda v^2(1 - \epsilon^2)(1 - R) , \\ Z_6 &= \frac{1}{2}\lambda(R - 1)\epsilon\sqrt{1 - \epsilon^2} , \end{aligned}$$


Approximate alignment without decoupling requires that  $|Z_6| \ll 1$  and  $\bar{m}^2 \sim \mathcal{O}(v^2)$ . To avoid  $\tan\beta$  very large or very small, we consider two limiting cases:  $|\epsilon| \ll 1$  and  $|R - 1| \ll 1$ .

In the limit of  $|\epsilon| \ll 1$ ,

$$m_h^2 = \frac{1}{2}\lambda v^2(1 + R) , \quad m_H^2 = \bar{m}^2 + \frac{1}{2}\lambda v^2(1 - R) , \quad c_{\beta-\alpha} = \frac{\lambda v^2(1 - R)\epsilon}{2(\bar{m}^2 - \lambda v^2 R)} .$$





In the limit of  $|R - 1| \ll 1$ ,

$$m_h^2 = \lambda v^2 , \quad m_H^2 = \bar{m}^2 , \quad c_{\beta-\alpha} = \frac{\lambda v^2(1 - R)\epsilon\sqrt{1 - \epsilon^2}}{2(\bar{m}^2 - \lambda v^2)} .$$



Allowed regions of the  $R$  vs.  $t_\beta \equiv \tan \beta$  parameter space for a Type-I (left) and Type-II (right) 2HDM with softly broken GCP symmetry, taking the precision  $h(125)$  LHC data into account.

We impose constraints from precision Higgs data, which favors a SM-like  $h(125)$ . We then chose  $m = 150$  GeV as a benchmark point. Together with the results for  $c_{\beta-\alpha}$  shown above, we find values of  $m_H$  and  $m_A$  of order 250 GeV.



## Regions of approximate alignment without decoupling

To be consistent with current LHC data, we shall also impose:


- Non-SM Higgs bosons in the regime of alignment without decoupling should have so far evaded LHC detection.
- Vector-like top quark bounds [we choose  $M_T \gtrsim 1.2$  TeV]
- Constraints on mixing between the top quark and its mirror partner<sup>5</sup> [ $\sin \theta_L \lesssim 0.12$ ]

The non-observation of non-SM Higgs bosons favors the low  $\tan \beta$  regime in Type-II models. The non-observation of vector-like quark effects in the top sector disfavors the regime of  $0.8 \lesssim \tan \beta \lesssim 2$ .

---

<sup>5</sup>See, e.g., A. Arhrib et al., Phys. Rev. D 97, 095015 (2018).

# LHC constraints on $H$ and $A$ masses in a Type-2 2HDM



Expected and observed 95% CL exclusion contour (left) in the MSSM  $m_h^{\text{mod+}}$  and (right) in the hMSSM scenarios. The expected median is shown as a dashed black line. The dark and bright gray bands indicate the 68 and 95% confidence intervals for the variation of the expected exclusion. The observed exclusion contour is indicated by the colored blue area. Taken from CMS Collaboration, JHEP **1809**, 007 (2018).

## Future work

- Adding in the mirror fermions corresponding to the down-type quarks and leptons.
- A detailed phenomenological study of the softly-broken GCP model to see the interplay between the spectrum of mirror fermions and the deviations from the alignment limit.
- Correlating the properties of the non-SM Higgs bosons with those of the mirror fermions.
- If mirror fermions are discovered, how to use data to identify the presence of an approximate GCP symmetry and to distinguish between GCP2 and GCP3.
- Assessing the extent of the fine-tuning of parameters in models of approximate alignment without decoupling (in the presence of an approximate symmetry).

Backup slides

## The alignment limit in equations

The CP-even Higgs squared-mass matrix yields,

$$Z_1 v^2 = m_h^2 s_{\beta-\alpha}^2 + m_H^2 c_{\beta-\alpha}^2,$$

$$Z_6 v^2 = (m_h^2 - m_H^2) s_{\beta-\alpha} c_{\beta-\alpha},$$

$$Z_5 v^2 = m_H^2 s_{\beta-\alpha}^2 + m_h^2 c_{\beta-\alpha}^2 - m_A^2.$$

If  $h$  is SM-like, then  $m_h^2 \simeq Z_1 v^2$  (i.e.,  $Z_1 \simeq 0.26$ ) and

$$|c_{\beta-\alpha}| = \frac{|Z_6| v^2}{\sqrt{(m_H^2 - m_h^2)(m_H^2 - Z_1 v^2)}} \simeq \frac{|Z_6| v^2}{m_H^2 - m_h^2} \ll 1,$$

If  $H$  is SM-like, then  $m_H^2 \simeq Z_1 v^2$  (i.e.,  $Z_1 \simeq 0.26$ ) and

$$|s_{\beta-\alpha}| = \frac{|Z_6| v^2}{\sqrt{(m_H^2 - m_h^2)(Z_1 v^2 - m_h^2)}} \simeq \frac{|Z_6| v^2}{m_H^2 - m_h^2} \ll 1.$$

# Family and Generalized CP symmetries of the 2HDM

## Higgs family symmetries

$$\mathbb{Z}_2 : \quad \Phi_1 \rightarrow \Phi_1, \quad \Phi_2 \rightarrow -\Phi_2$$

$$\Pi_2 : \quad \Phi_1 \longleftrightarrow \Phi_2$$

$$U(1)_{\text{PQ}} \text{ [Peccei-Quinn]}: \quad \Phi_1 \rightarrow e^{-i\theta}\Phi_1, \quad \Phi_2 \rightarrow e^{i\theta}\Phi_2$$

$$SO(3): \quad \Phi_a \rightarrow U_{ab}\Phi_b, \quad U \in U(2)/U(1)_Y$$

## Generalized CP (GCP) transformations

$$\text{CP1} : \quad \Phi_1 \rightarrow \Phi_1^*, \quad \Phi_2 \rightarrow \Phi_2^*$$

$$\text{CP2} : \quad \Phi_1 \rightarrow \Phi_2^*, \quad \Phi_2 \rightarrow -\Phi_1^*$$

$$\text{CP3} : \quad \Phi_1 \rightarrow \Phi_1^*c_\theta + \Phi_2^*s_\theta, \quad \Phi_2 \rightarrow -\Phi_1^*s_\theta + \Phi_2^*c_\theta, \quad \text{for } 0 < \theta < \frac{1}{2}\pi$$

where  $c_\theta \equiv \cos \theta$  and  $s_\theta \equiv \sin \theta$ .